



Published in final edited form as:

Am J Gastroenterol. 2015 February ; 110(2): 223–263. doi:10.1038/ajg.2014.435.

ACG Clinical Guideline: Genetic Testing and Management of Hereditary Gastrointestinal Cancer Syndromes

Sapna Syngal, MD, MPH, FACG^{1,2,3}, Randall E. Brand, MD, FACG⁴, James M. Church, MD, FACG^{5,6,7}, Francis M. Giardiello, MD⁸, Heather L. Hampel, MS, CGC⁹, and Randall W. Burt, MD, FACG¹⁰

¹ Brigham and Women's Hospital, Boston, Massachusetts, USA

² Dana Farber Cancer Institute, Boston, Massachusetts, USA

³ Harvard Medical School, Boston, Massachusetts, USA

⁴ Department of Medicine, Division of Gastroenterology, Hepatology and Nutrition, University of Pittsburgh Medical Center, Pittsburgh, Pennsylvania, USA

⁵ Department of Colorectal Surgery, Cleveland Clinic, Cleveland, Ohio, USA

⁶ Sanford R Weiss, MD, Center for Hereditary Colorectal Neoplasia, Cleveland Clinic Foundation, Cleveland, Ohio, USA

⁷ Digestive Disease Institute, Cleveland Clinic Foundation, Cleveland, Ohio, USA

⁸ Johns Hopkins University School of Medicine, Baltimore, Maryland, USA

⁹ Department of Internal Medicine, Ohio State University, Columbus, Ohio, USA

¹⁰ Huntsman Cancer Institute, University of Utah School of Medicine, Salt Lake City, Utah, USA.

Abstract

This guideline presents recommendations for the management of patients with hereditary gastrointestinal cancer syndromes. The initial assessment is the collection of a family history of cancers and premalignant gastrointestinal conditions and should provide enough information to develop a preliminary determination of the risk of a familial predisposition to cancer. Age at diagnosis and lineage (maternal and/or paternal) should be documented for all diagnoses, especially in first- and second-degree relatives. When indicated, genetic testing for a germline

Correspondence: Sapna Syngal, MD, MPH, FACG, Dana Farber Cancer Institute, 450 Brookline Avenue, Dana 1124, Boston, Massachusetts 02215, USA. sapna_syngal@dfci.harvard.edu.

CONFLICT OF INTEREST

Guarantor of the article : Sapna Syngal, MD, MPH, FACG.

Specific author contributions: Each of the authors (Syngal, Hampel, Burt, Brand, Giardiello, and Church) played a role in writing portions of the manuscript. Dr Syngal assimilated the sections into the comprehensive document and each co-author subsequently provided feedback in conference calls and written form on subsequent revisions.

Potential competing interests : Dr Syngal has served as a consultant to EXACT Sciences and Merck. She is an unpaid collaborator/advisor to Myriad Genetics. One of her mentees receives research grant support from Myriad Genetics. Dr Syngal does not receive salary support from this grant, but is a collaborator on the project. Dr Brand has served as a consult to Myriad Genetics and is a Scientific Advisory Board Member for Asuragen. Ms Hampel has received honoraria from Quest Diagnostics and InVita Laboratory for educational talks in the past year. In addition, she receives research genetic testing support from Myriad Genetics. Dr Burt consults for Myriad Genetics. The authors have no relationships with the funding sources specified.

mutation should be done on the most informative candidate(s) identified through the family history evaluation and/or tumor analysis to confirm a diagnosis and allow for predictive testing of at-risk relatives. Genetic testing should be conducted in the context of pre- and post-test genetic counseling to ensure the patient's informed decision making. Patients who meet clinical criteria for a syndrome as well as those with identified pathogenic germline mutations should receive appropriate surveillance measures in order to minimize their overall risk of developing syndrome-specific cancers. This guideline specifically discusses genetic testing and management of Lynch syndrome, familial adenomatous polyposis (FAP), attenuated familial adenomatous polyposis (AFAP), *MUTYH*-associated polyposis (MAP), Peutz–Jeghers syndrome, juvenile polyposis syndrome, Cowden syndrome, serrated (hyperplastic) polyposis syndrome, hereditary pancreatic cancer, and hereditary gastric cancer.

Hereditary gastrointestinal (GI) cancer syndromes represent a phenotypically diverse group of disorders that exhibit distinct patterns of inheritance in an individual's progeny. Over the past few decades, the expansion of familial cancer registries and advancement in genomics have led to the development of clinical diagnostic criteria for specific hereditary syndromes as well as the discovery of multiple genes in which germline mutations predispose individuals to syndrome-associated neoplastic manifestations. This guideline first discusses essential elements of a patient's personal and family history that allow for risk assessment for potential inherited cancer susceptibility. It then addresses the currently most well-characterized GI cancer susceptibility syndromes: Lynch syndrome (LS), familial adenomatous polyposis (FAP), attenuated familial adenomatous polyposis (AFAP), *MUTYH*-associated polyposis (MAP), Peutz–Jeghers syndrome (PJS), juvenile polyposis syndrome (JPS), Cowden syndrome (CS), serrated (hyperplastic) polyposis syndrome, hereditary pancreatic cancer, and hereditary gastric cancer. For each of these syndromes, we outline diagnostic criteria and indications for genetic evaluation, describe the currently known associated underlying genes, and make recommendations for surveillance and management of at-risk individuals and those found to carry a definitive disease-causing mutation. Finally, we discuss the elements of informed consent that must accompany genetic evaluation as well as currently evolving genetic testing technologies that may change how genetic testing is conducted in the near-term future.

Each section of the document presents summary statements, the key recommendations related to the section topic, followed by a summary of the supporting evidence (**Tables 1 and 2**). A search of MEDLINE via the OVID interface using the MeSH term “hereditary cancer syndrome” limited to clinical trials, reviews, guidelines, and meta-analysis for the years 1966–2013 was performed to develop the document and create summary statements and recommendations. “Summary statements” and “recommendations” are distinguished by whether it was possible to address the quality of evidence supporting the statements based on an objective grading system. An objective measure that provides assessment of the strength of data regarding prognostic indicators does not currently exist, and similarly, “motherhood” statements (such as the importance of obtaining a family history) that are based on sound clinical judgment are often not subject to systematic clinical studies as they are understood to reflect sound clinical practice. The summary statements therefore reflect consensus opinion by the authors and a thorough literature review that reflects expert

opinion by leaders in the field and other consensus guidelines. For management recommendations, where alternative strategies are and should be subject to rigorous assessment, the GRADE (Grading of Recommendations Assessment, Development and Evaluation) system was used to grade the strength of recommendations and the quality of evidence (1). An explanation of the quality of evidence and strength of recommendations is shown in **Table 3**. The quality of evidence, which influences the strength of the recommendation, ranges from “high” (further research is very unlikely to change our confidence in the estimate of effect) to “moderate” (further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate) to “low” (further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate), and to “very low” (any estimate of effect is uncertain). The strength of a recommendation is graded as strong when the desirable effects of an intervention clearly outweigh the undesirable effects and is graded as conditional when uncertainty exists about the trade-offs.

The field of cancer genetics poses some challenges with respect to the GRADE system. Because of the rarity of the syndromes, and the relatively recent discovery of cancer susceptibility genes, data regarding long-term outcomes regarding optimal management strategies at this time are limited to observational studies. Randomized clinical trials, which are the gold standard of systems such as GRADE, are difficult to conduct in rare diseases, where the main objective outcome, reduction in cancer mortality, takes years to assess and large patient numbers. The reader, therefore, should take the assessments of quality of evidence with caution— the often “low” or “very low” quality gradings reflect primarily a lack of available data and not that the quality of studies conducted thus far has been poor.

STANDARDS FOR MINIMAL CANCER FAMILY HISTORY ASSESSMENT IN GI PRACTICE

Summary statements

- A family history of cancer and premalignant GI conditions that provides sufficient information to develop a preliminary determination of the risk of a familial predisposition to cancer should be obtained for all patients being evaluated in outpatient gastroenterology and endoscopy practices.
- Essential elements of a family history include presence and type of cancer diagnoses in first- and second-degree relatives, and presence and (ideally) type of polyps in first-degree relatives; age and lineage should be noted for each diagnosis.

Summary of evidence

Approximately 5–10% of cancers are attributable to a hereditary cancer predisposition syndrome. Identifying those patients who have an inherited cancer predisposition syndrome has significant benefit to both the patient and at-risk relatives. For the index patient, the diagnosis of a hereditary cancer syndrome has implications for his/her surveillance strategy for multiple component tumors in terms of age of initiation and intervals between surveillance exams, and may lead to the consideration of prophylactic surgery or more

extensive surgery in the case of neoplasia development. the diagnosis of an inherited syndrome also has significant implications for management of the patient's immediate and extended family.

Features of a patient's personal history may be the initial clue to the possibility of an inherited predisposition to cancer. Hallmark features, whose specifics are outlined in detail in the remainder of this guideline, include early age at onset of polyps or cancer and unusual numbers or histologies of cancers or premalignant conditions. Family history is the other key component to the identification of those individuals who may have an inherited predisposition to malignancy or who are at increased risk for additional primary cancers.

The goal of any cancer family history, in combination with the patient's personal history, is to provide enough information to make a preliminary determination about whether the patient may have a familial predisposition to cancer, may benefit from genetic counseling and possibly testing for underlying cancer susceptibility genes, or may not need genetic counseling and testing, but still require more intensive surveillance than the average- or moderate-risk patient. As there is currently no clear evidence base to define how family history should be taken or what constitutes the right amount of information for an initial cancer screening family history, an expert panel was recently convened to define how and what to collect for a family cancer history (2). The panel agreed that although the gold standard family history is the comprehensive, three-generation pedigree used in medical genetics, counseling, and research settings, this evaluation is time consuming and not feasible in general medical practice. For most patients, family history of cancer and premalignant conditions in close relatives is most relevant. Guidelines for consideration of genetic risk assessment, such as the NCCN Clinical Guidelines in Oncology for Genetic/Familial High-Risk Assessment: Breast and Ovarian, (3) focus on first- and second-degree relatives, although they may optionally incorporate family history in third-degree relatives. Thus, family history of cancer in first-degree (parents, children, and siblings) and second-degree (grandparents, aunts/uncles, nieces/nephews, grandchildren, and half-siblings) relatives is often sufficient to assess a patient's empiric risk of common cancers or a cancer patient's risk of a second primary cancer. Relatives' age at cancer or polyp diagnosis should also be assessed because this factors into both genetic risk assessment guidelines and cancer screening recommendations. Maternal and paternal lineages should be assessed separately. Accuracy of self-reported cancer family history in first-degree relatives (FDRs) has been shown to be >75% for most cancers, including colorectal, breast, ovarian, and pancreatic cancers. Studies have shown a decrease in the accuracy of reported family history in more distant relatives, ranging from 50 to 80% depending on the cancer (4,5). Hence, the routine review of family medical records, although not required during family history collection, can be helpful in particular cases where the cancer site is in question.

LYNCH SYNDROME (LS)

Tumor testing and indications for genetic testing

Summary statements

1. All newly diagnosed colorectal cancers (CRCs) should be evaluated for mismatch repair deficiency.
2. Analysis may be done by immunohistochemical testing for the *MLH1/MSH2/MSH6/PMS2* proteins and/or testing for microsatellite instability (MSI). Tumors that demonstrate loss of *MLH1* should undergo BRAF testing or analysis for *MLH1* promoter hypermethylation.
3. Individuals who have a personal history of a tumor showing evidence of mismatch repair deficiency (and no demonstrated BRAF mutation or hypermethylation of *MLH1*), a known family mutation associated with LS, or a risk of 5% chance of LS based on risk prediction models should undergo genetic evaluation for LS.

Summary of evidence

LS, the most common cause of inherited CRC, is an autosomal-dominant condition defined by the presence of a germline mutation in a DNA mismatch repair gene (or *EPCAM*). It was often previously referred to as hereditary nonpolyposis colorectal cancer. LS tumors are associated with changes in the length of nucleotide repeat sequences of tumor DNA, termed MSI. MSI results from defective mismatch repair and is associated with loss of expression of the *MLH1*, *MSH2*, *MSH6*, and/or *PMS2* proteins that can be detected by immunohistochemical (IHC) analysis. Multiple international studies have demonstrated that the prevalence of MSI in population-based series of CRC ranges from 7 to 19% (6–10). The sensitivity of MSI testing among those with *MLH1* or *MSH2* mutations is 80–91%, and is 55–77% among those with *MSH6* or *PMS2* mutations; the specificity of MSI testing is 90% (11). The sensitivity of IHC testing, regardless of the MMR gene involved, is 83% and the specificity is 89% (11). MSI and IHC results are highly correlated (9,12), and as protein staining is often easier to perform than DNA analysis in a clinical setting, it may be a more feasible option for widespread MSI screening. In order to facilitate surgical planning, tumor testing on suspected CRC should be performed on preoperative biopsy specimens if possible. For individuals whose IHC indicates loss of the *MLH1* protein, determination of the mechanism of loss should be pursued as an additional screening step, and this may be done by analysis for a BRAF mutation or promoter hypermethylation studies. Almost no LS tumors carry a BRAF mutation, whereas 68% of those without LS do (11). Individuals who demonstrate evidence of MMR deficiency, independent of somatic *MLH1* silencing, should undergo genetic testing.

Genetic etiology

Summary statement

- Genetic testing of patients with suspected LS should include germline mutation genetic testing for the *MLH1*, *MSH2*, *MSH6*, *PMS2*, and/or *EPCAM* genes (13–23), or the altered gene(s) indicated by IHC testing.

Summary of evidence

In 1993, genome-wide linkage analysis in several large families with autosomal-dominant CRC and the demonstration of associated tumor MSI led to the subsequent cloning of the

mismatch repair genes *MLH1* and *MSH2*, followed by *MSH6*, *PMS2*, and *EPCAM* (13–23). Multiple large international population-based series have demonstrated that MMR gene mutations account for 1–3% of newly diagnosed CRC cases (6,7,9,10,12). LS should be considered in individuals whose tumors show evidence of MMR deficiency as discussed above (without the presence of a *BRAF* mutation or *MLH1* promoter hypermethylation), and those whose personal and/or family history fulfill the Amsterdam criteria, Bethesda Guidelines, or who have a 5% risk of carrying a germline mutation based on available prediction models (24,25) (**Table 4**). The computational models all appear to outperform existing clinical guidelines (25), primarily because of limited sensitivity of the clinical criteria in identifying mutation carriers. In families where LS is a consideration, and no tumor sample is available for analysis, direct germline testing of an unaffected at-risk individual whose risk is calculated to be 5% based on the PREMM1,2,6 risk prediction model (accessible at <http://premm.dfci.harvard.edu/>) is a strategy that has been demonstrated to be cost effective in improving health outcomes (26). The complexity of clinical criteria may be difficult to apply in clinical practice. A simple, validated three-question tool may be used as a quick initial screen in busy practices to identify which patients need further risk assessment (27) (**Table 4**).

Surveillance and management of CRC

Recommendation

1. In individuals at risk for or affected with LS, screening for CRC by colonoscopy should be performed at least every 2 years, beginning between ages 20 and 25 years. Annual colonoscopy should be considered in confirmed mutation carriers (strong recommendation, moderate quality of evidence for screening, and very low quality of evidence for annual surveillance and age of initiation).

Summary of evidence

The precursor lesion for a LS-related CRC is an adenomatous polyp that is often proximal and can occasionally be flat rather than elevated/polypoid and likely to demonstrate villous features, high-grade dysplasia, and a preponderance of tumor-infiltrating lymphocytes (28). The historical term nonpolyposis CRC was intended to differentiate this condition from FAP in which patients develop hundreds of adenomas. LS patients develop few (usually <10) early-onset adenomas, but the adenoma–carcinoma sequence appears to be accelerated in LS with polyp to cancer dwell times estimated at 35 months compared with 10–15 years in sporadic cancer (29). Reports of lifetime risks of CRC for *MLH1* and *MSH2* gene mutation carriers range from 22 to 74% (30–35) (**Table 5**). Lower risk for colorectal malignancy has been found in women, but not in men with *MSH6* mutations (30% vs. 69% cumulative risk by age 70 years, respectively), compared with *MLH1* and *MSH2* carriers (36). CRC risk is reported to be lower in one study of 99 *PMS2* mutation carriers, with an estimated cumulative risk of 15–20% by age 70 years. (37). The mean age of CRC diagnosis in LS patients is 44–61 years (8,12,38,39) compared with 69 years in sporadic cases of CRC (40).

CRC prevention in LS families is guided by the distinctive characteristics of these malignancies, including the younger age of presentation, right-sided colorectal

predominance, and rapid polyp growth with shorter dwell time before malignant conversion. Evidence for the effectiveness of colorectal screening in decreasing CRC mortality has been documented in studies by Järvinen *et al.* (41–43) (**Table 6**). Individuals at risk for LS who took up colonoscopic surveillance had 65% ($P=0.003$) less death from CRC compared with those who refused surveillance. Update of this Finnish study that analyzed colonoscopic surveillance in LS mutation carriers found no difference in CRC death between mutation carriers and mutation-negative relatives (43). Dove-Edwin *et al.* (44) reported the results of a prospective observational study of colonoscopy surveillance of members in hereditary nonpolyposis colorectal cancer or LS families, revealing a 72% decrease in mortality from CRC in those undergoing screening. In several studies (45–48), more frequent colonoscopy screening (2 years) was associated with an earlier stage of CRC at diagnosis and less CRC than less frequent colonoscopy. At least every 2-year colonoscopic surveillance of LS patients is supported by the data above and the rapid adenoma–carcinoma sequence reported in these patients. In surveillance of MMR germline mutation-positive patients, consideration should be given to annual colonoscopy, as several studies have demonstrated CRC development with surveillance intervals that are between 1 and 2 years (29,47). In carriers of deleterious *MSH6* and *PMS2* mutations, the risk of CRC is less and age of diagnosis later (37,49) than in patients with *MLH1* and *MSH2* mutations. In these affected individuals, consideration could be given to starting surveillance at age 25–30 in *MSH6* and *PMS2* carriers (24), unless an early-onset cancer exists in a given family; however, this approach of gene-specific alterations in surveillance has not been evaluated in clinical studies.

Recommendation

2. Colectomy with ileorectal anastomosis (IRA) is the preferred treatment of patients affected with LS with colon cancer or colonic neoplasia not controllable by endoscopy. Segmental colectomy is an option in patients unsuitable for total colectomy if regular postoperative surveillance is conducted (conditional recommendation, moderate quality of evidence).

Summary of evidence

The treatment for patients with CRC or premalignant polyps that cannot be removed by colonoscopy is subtotal colectomy with IRA. A high rate of metachronous CRC (16% at 10 years; 41% at 20 years) is noted in LS patients who have undergone segmental surgical resection of the initial CRC in several retrospective studies (48,50,51). A standard low anterior resection or abdominal perineal resection may be performed to treat rectal cancers in LS patients, although the residual colon is at high risk of metachronous neoplasia. Younger patients may be offered a total proctocolectomy and ileal pouch anal anastomosis (IPAA), or ileostomy. A recent retrospective study of 79 LS patients with rectal cancer who had undergone proctectomy found a cumulative risk of metachronous colon cancer to be 19% at 10 years, 47% at 20 years, and 69% at 30 years after surgical resection (51). The risk of metachronous cancer is substantially abated if extensive colectomy is performed (0–3.4%) (48,50,51).

In a Dutch study, no difference in global quality of life was noted between 51 LS patients who underwent partial colectomy and 53 patients who underwent subtotal colectomy,

although functional outcome (stool frequency, stool-related aspects, social impact) was worse after subtotal colectomy than after partial colectomy (52). A comparison of life expectancy gained performing total colectomy vs. hemicolectomy in LS patients at ages 27, 47, and 67 years by Markov modeling was 2.3, 1, and 0.3 years, respectively. These investigators concluded that total colectomy is the preferred treatment in LS but hemicolectomy may be an option in older individuals. Consideration for less extensive surgery may be given in patients who are >60–65 years of age.

The option of prophylactic colectomy should be discussed with mutation carriers who have an endoscopically normal colon as an alternative to surveillance. Although rarely chosen, it may be attractive to patients from families where the prevalence of colon cancer is very high, or for whom colonoscopy is difficult. Direct comparative studies of extensive surgery vs. annual or biennial surveillance have not been conducted, and are unlikely to be instituted because of the multitude of patient-dependent factors that affect comorbidities and quality of life.

Surveillance and management of extracolonic malignancies

Gynecologic malignancies

Recommendations

3. Hysterectomy and bilateral salpingo-oophorectomy should be offered to women who are known LS mutation carriers and who have finished child bearing, optimally at age 40–45 years (conditional recommendation, low quality of evidence).
4. Screening for endometrial cancer (EC) and ovarian cancer should be offered to women at risk for or affected with LS by endometrial biopsy and transvaginal ultrasound annually, starting at age 30 to 35 years before undergoing surgery or if surgery is deferred (conditional recommendation, very low quality of evidence).

Summary of evidence

EC is the second most common cancer occurring in LS. Estimates of the cumulative lifetime risk of EC in LS patients range from 15 to 71%, with variability depending on specific gene mutation (31,33,35–37,49) (**Table 7**); reports of age at diagnosis of this malignancy are clearly a decade or more younger than sporadic EC but range from 48 to 54 years (31,33,35–37,49) (**Table 7**). Estimates of the cumulative lifetime risk of ovarian cancer in LS patients range from 3.4 to 22% (31,38,53–55) (**Table 7**).

Because of the worrisome cumulative risk of EC, several annual screening modalities have been proposed including pelvic exams, transvaginal ultrasound, endometrial sampling, and CA 125 testing. Few studies of these interventions have been conducted and there is currently no evidence of survival benefit from EC surveillance (**Table 8**). Decrease in death from EC screening may be difficult to prove as 75% of LS patients with EC present with stage 1 disease and have an 88% 5-year survival rate. Transvaginal ultrasound has poor sensitivity and specificity for the diagnosis of EC in this population (56,57). However,

endometrial sampling appears useful in identifying some asymptomatic patients with EC and those with premalignant endometrial lesions (**Table 8**). Currently, no studies on the effectiveness of ovarian screening are available for women in LS families.

One retrospective study of 315 women with MMR mutations who did and did not have hysterectomy and oophorectomy revealed no cancers in the surgical group compared with a 33% and 5.5% rate of uterine and ovarian cancer, respectively, in the nonsurgical group (58). Cost-effectiveness analysis modeling of gynecological screening vs. prophylactic gynecological surgery (hysterectomy and bilateral salpingo-oophorectomy) in a theoretical population of 30-year-old women with LS revealed that prophylactic surgery had lower cost and higher quality-adjusted life-years (59). An additional modeling study evaluated multiple screening and surgical strategies. This investigation concluded that annual screening starting at age 30 years followed by prophylactic surgery at age 40 years was the most effective gynecologic cancer prevention strategy, but incremental benefit over prophylactic surgery at age 40 years alone was attained at substantial cost (60).

Recommendations

5. Screening for gastric and duodenal cancer can be considered in individuals at risk for or affected with LS by baseline esophagogastroduodenoscopy (EGD) with gastric biopsy at age 30–35 years, and treatment of *Helicobacter pylori* infection when found. Data for ongoing regular surveillance are limited, but ongoing surveillance every 3–5 years may be considered if there is a family history of gastric or duodenal cancer (conditional recommendation, very low quality of evidence).
6. Screening beyond population-based recommendations for cancers of the urinary tract, pancreas, prostate, and breast is not recommended unless there is a family history of the specific cancers (conditional recommendation, low quality of evidence).

Summary of evidence

The impact of a family history of extracolonic cancers on other at-risk relatives has not been systematically studied. Some studies show clustering of extracolonic cancers in families, whereas others have not (discussed when available in section below for individual cancers). In clinical practice, decision making regarding surveillance for extracolonic cancers is generally done on a case-by-case basis, taking into account cancer history in at-risk first- and second-degree relatives on the affected side of the family.

Some studies have estimated the lifetime risk of gastric cancer in LS to be as high as 13%, but it is currently much lower in North America and Western Europe. A carefully conducted time trend study of gastric cancer found an 8.0% and 5.3% lifetime risk of this malignancy in males and females with MMR gene mutation, respectively, and lack of familial clustering (61). The majority of gastric cancers in LS patients appear to be histologically classified as intestinal type (61,62), and, consequently, potentially amenable to endoscopic surveillance. There are no studies that have evaluated the effectiveness of screening and surveillance for gastric cancer in LS patients.

The lifetime risk for small bowel cancer ranges from 0.4 to 12.0% (31,34,53,54,63,64). The majority of small bowel cancers in a LS cohort were located in the duodenum or ileum (65) and within the reach of EGD and colonoscopy with dedicated ileal intubation. Studies of small bowel screening in LS patients are lacking. However, one screening investigation of 35 gene mutation carriers found that 2 had jejunal adenomas and 1 had a jejunal cancer (66) (**Table 8**). Six additional patients had capsule endoscopy images of uncertain clinic relevance, prompting further invasive investigation in five patients. A recent publication suggested that routine surveillance of the small bowel in LS was not cost efficient (55).

Estimates of the lifetime risk of urinary tract cancer in LS range from 0.2 to 25%, depending on the study and which urinary tract cancers are included (transitional cell carcinoma of the ureter, renal pelvis, and bladder) (31,53,54,63,64,67). Currently, a dearth of literature on screening for urinary cancer in LS patients exists. One retrospective study evaluating screening for urinary cancer by urine cytology in individuals in hereditary nonpolyposis colorectal cancer or LS families found a poor (29%) sensitivity in diagnosing cancer in asymptomatic patients and production of many false positive results requiring invasive investigation (68) (**Table 8**). Screening has not been shown to be effective with urine cytology and urinalysis for microscopic hematuria for urinary cancer in the general population and in groups at higher risk for bladder cancer from environmental factors (69,70). The benefit of ultrasound screening is unknown. In summary, limited data exist to advocate urinary screening.

The risk of pancreatic cancer (PC) in LS patients was noted to be elevated in two cohort studies. In one study, the standardized incidence ratio for PC was 10.7 (95% confidence interval, 2.7–47.7), with a 10-year cumulative risk of 0.95% (71), and the other study reported a 8.6-fold increase (95% confidence interval, 4.7–15.7), with cumulative risk of 3.7% by age 70 years (72). The benefit of screening for PC in LS has not been evaluated. An international pancreas consensus panel recommended that, based on expert opinion, mismatch repair gene mutation carriers with one affected FDR should be considered for annual PC surveillance with magnetic resonance imaging (MRI) and/or endoscopic ultrasound based on early data in other cohorts with comparable risk (73).

There are conflicting data regarding the risk of several other extracolonic cancers in patients with LS. The relationship between LS and breast cancer is unclear. Although a small increase in lifetime breast cancer risk of 18% has been found (54,74), most clinic-based registry reports have not demonstrated this consistently (55,75). In two recent studies the relative risk of prostate cancer is 2.0- to 2.5-fold the general population risk (54,76); however, the effectiveness of intensive screening beyond population recommendations has not been evaluated.

Prevention strategies

Diet, exercise, smoking, and supplements—A prospective analysis of 386 patients with LS undergoing surveillance revealed that current smokers had an increased risk of colorectal adenomas compared with past smokers and never smokers (hazard ratio of 6.1 vs. 3.0 vs. 1, respectively (77). Excess body weight (body mass index >25 kg/m²) has been shown to be associated with an elevated risk (hazard ratio of 8.7 compared with normal

weight) of colorectal adenomas in men with LS in the same cohort; an elevated risk was not found in women with a high body mass index (78).

Chemoprevention

Recommendation

7. Although data suggest that daily aspirin may decrease the risk of colorectal and extracolonic cancer in LS, currently the evidence is not sufficiently robust or mature to make a recommendation for its standard use (conditional recommendation, moderate quality of evidence).

Summary of evidence

Resistant starch and aspirin have been assessed as chemopreventive agents in patients with LS. The Colorectal/Adenoma/Carcinoma Prevention Programme 2 (CAPP2) was a randomized placebo-controlled trial with a two-by-two design investigating the effect of resistant starch (Novelose) 30 g per day and aspirin 600 mg per day taken for up to 4 years on development of colorectal adenoma and cancer (79). This study randomized 727 participants to starch or placebo and 693 between aspirin and placebo. The use of resistant starch, aspirin, or both had no effect on the incidence of colorectal neoplasia in LS carriers over a mean follow-up period of 29 months. The CAPP2 investigators subsequently evaluated the long-term effect of 600 mg of aspirin usage on CRC development (80). At a mean follow-up of 55.7 months, intention-to-treat analysis of time to first CRC showed a hazard ratio of 0.63 (95% confidence interval, 0.35–1.13, $P=0.12$). An intention-to-treat analysis of all LS cancers (colorectal, endometrial, ovarian, pancreatic, small bowel, gallbladder, ureter, stomach, kidney, and brain) revealed a protective effect of aspirin vs. placebo (hazard ratio, 0.65; 95% confidence interval, 0.42–1.00, $P=0.05$). During the intervention, adverse events did not differ between aspirin and placebo groups.

The CAPP2 trial has several limitations. First, ascertainment of the end point, CRC, was not standardized, and more intensive colonoscopic evaluation could have occurred in the aspirin group than in the non-aspirin group because of more frequent adverse effects after intervention. Second, the extracolonic cancers did not undergo molecular evaluation to assess whether they were related to the germline MMR mutation. In addition, the dose of daily aspirin utilized in the CAPP2 trial is significantly higher than that noted to be effective (75 mg a day) in sporadic CRC chemoprevention.

The CAPP3 trial is currently underway to establish the optimum dose and duration of aspirin treatment. Although data exist to suggest that aspirin may decrease the risk of colorectal and extracolonic cancer in LS, currently the evidence is not sufficiently robust or mature to make a recommendation for its standard use.

Adenomatous polyposis syndromes

Familial adenomatous polyposis/*MUTYH*-associated polyposis/attenuated polyposis indications for genetic evaluation—*Summary statement*

- Individuals who have a personal history of >10 cumulative colorectal adenomas, a family history of one of the adenomatous polyposis syndromes, or a history of adenomas and FAP-type extracolonic manifestations (duodenal/ampullary adenomas, desmoid tumors (abdominal>peripheral), papillary thyroid cancer, congenital hypertrophy of the retinal pigment epithelium, epidermal cysts, osteomas) should undergo assessment for the adenomatous polyposis syndromes.

Summary of evidence

There are three known hereditary syndromes where inheritance of germline mutations produces enhanced colorectal carcinogenesis, manifested by early age of onset of multiple colorectal adenomas with the potential for early development of CRC: FAP, AFAP and MAP.

FAP is defined by the presence of ≥100 synchronous colorectal adenomas inherited in an autosomal-dominant manner. Estimates of the prevalence of FAP vary from 1 in 6,850 to 1 in 31,250 live births (2.29 to 3.2 cases per 100,000 individuals) (81–87). The frequency is fairly constant throughout the world, with men and women being affected equally.

Patients with 10 to 99 synchronous adenomas have oligopolyposis or AFAP. AFAP is defined by <100 adenomas at presentation inherited in an autosomal-dominant pattern. Patients with AFAP have fewer adenomas than those with typical FAP, averaging 25 polyps in one study and exhibiting a more proximal colonic preponderance than in typical FAP (88–90). However, the polyp number is extremely variable within many kindreds.

In 2002, an attenuated polyposis syndrome was described in three siblings affected with multiple adenomas and/or CRC inherited in an autosomal-recessive pattern (91). This recessive condition is referred to as MAP and is characterized by an increased risk for CRC and multiple adenomatous polyps that can mimic FAP or AFAP.

There are two settings in which to consider genetic testing for the adenomatous polyposis syndromes: (i) testing an individual with a phenotype suggestive of one of the polyposes, but where the clinical diagnosis is not certain, and (ii) testing relatives of a patient with a known germline mutation. The first setting is usually defined as a patient with ≥10 cumulative adenomas, or sometimes suggestive extracolonic manifestations, but no known family history of an underlying pathogenic mutation. Genetic testing in this setting should be comprehensive and the absence of a mutation does not definitively rule out a clinical diagnosis if the phenotype is striking. In the second setting, relatives of an individual with a known pathogenic mutation are tested for the presence or absence of that particular mutation. A positive test indicates the diagnosis of a syndrome, whereas a negative test (absence of mutation) rules it out and establishes that the individual is not at a syndromic risk for cancer and polyps.

Genetic etiology

Summary statement

- Genetic testing of patients with suspected adenomatous polyposis syndromes should include *APC* and *MUTYH* gene mutation analysis.

Summary of evidence

FAP arises from germline mutations of the *APC* gene on chromosome 5q21 (92,93). FAP is dominantly inherited and is close to 100% penetrant. Up to one-third of newly diagnosed cases not belonging to previously identified families appear to represent either *de novo* germline mutations or mosaicism (81,94–98).

APC is a tumor suppressor gene; thus, gene inactivation occurs only after both alleles are mutationally damaged. In FAP, one allele is inherited in a mutated form. Adenoma formation is initiated when the second allele is damaged or lost by a somatic event. The progression of adenoma to carcinoma after *APC* inactivation is similar in FAP and the sporadic setting in that mutations accumulate in additional relevant genes including *K-ras*, *p53*, a gene or genes on chromosome 18, and possibly others (99). Although mutations have been found scattered throughout the *APC* gene, most are located in the 5' end of exon 15, often called the *mutation cluster region* (100,101). The location of mutations in the *APC* gene correlate to some degree with colonic adenoma number, desmoid tumor occurrence, and congenital hypertrophy of the retinal pigment epithelium (102). Individuals with >1,000 polyps exhibit mutations in the mid-portion of the gene (102,103).

AFAP arises from *APC* mutations at either the far proximal (5') end of the gene, the far distal (3') end of the gene, or in certain locations of exon 9 (90,102). Whole or partial gene deletions may also give an attenuated phenotype.

MAP is a recessively inherited syndrome due to biallelic (homozygous or compound heterozygous) *MUTYH* mutations. *MUTYH* is a base excision repair gene involved in DNA oxidative damage repair whose protein repairs oxidative damage to the DNA (91). Failure of base excision repair results in CG–AT transversions in multiple genes, including *APC* and *KRAS* (91,104,105). Polyp and cancer predisposition occur with germline *MUTYH* mutations, but somatic *MUTYH* mutations do not appear to play a role in the pathogenesis of colon cancer (106,107). The two most prevalent *MUTYH* mutations, occurring in >80% of individuals of European ancestry with MAP, are two missense mutations Y179C and G396D (previously referred to as Y165C and G382D, respectively) (91,104,106,108–110). Other population-specific *MUTYH* mutations have been found (107). Approximately 90% of “western” population MAP patients have at least one of these two mutations; however, many other distinct *MUTYH* mutations have been reported (107).

MAP is most commonly found in patients presenting with 20 to 99 adenomas (107,111). Biallelic *MUTYH* mutations are found in 7.5% to 12.5% of patients with >100 adenomas in whom a disease-causing *APC* mutation is not found (104,112) and in 16 to 40% of patients with 15 to 99 colonic adenomas but not FAP (104,108,109,113,114). Although biallelic mutations have been found in individuals with early-onset CRC and few to no polyps, and in individuals with <10 adenomas without CRC, this is relatively uncommon (115). MAP was found in 0 out of 400 individuals with <4 adenomas, 2 of 444 (0.5%) unselected CRCs, and 0 out of 62 MSI high CRCs (112). Similar frequencies among patients with polyps and

CRCs have been found in other studies. In a study from Finland, 0.4% of 1,042 population-based CRC cases had biallelic *MUTYH* mutations (110). All those with mutations were found to have adenomas, ranging from 3 to 100. In a large (2,239 cases and 1,845 controls) population-based case-control study from Scotland, 0.8% of CRC cases <55 years old and 0.54% of all cases had biallelic mutations (116).

Monoallelic *MUTYH* mutations are found in 1 to 2% of the general population (107). Monoallelic *MUTYH* mutation carriers may have a slightly elevated risk of CRC, although the precise magnitude of the increased risk is currently unclear; most studies estimate a 1.5–2-fold risk above the general population (107). There is currently no consensus regarding the management of monoallelic carriers as data are limited. An option for clinicians at the current time is to manage monoallelics as individuals with a FDR with CRC, offering colonoscopy as a surveillance modality every 5 years, beginning 10 years earlier than the earliest CRC diagnosis.

Polymerase-proofreading associated polyposis is a newly described syndrome and only a few families have been characterized (117). Its phenotype includes oligo-adenomatous polyposis and an early age of onset of colorectal and EC. In a recent study of 858 familial/early-onset CRC cases and polyposis, one known *POLE* germline mutation and one new *POLD1* mutation were identified (118). Polymerase proofreading-associated polyposis is dominantly inherited and penetrance appears high.

Patients with clinical suspicion of an adenomatous polyposis syndrome should have genetic counseling and testing for germline mutations in *APC* and *MUTYH*. Failure to identify a mutation in an index case does not rule out the diagnosis of adenomatous polyposis, as mutations cannot be found in all families. If testing is negative, and clinical suspicion remains high, testing for other possible underlying genes should be considered. Failure to find a mutation means that all close relatives must still be screened as if they have FAP.

Finding a mutation confirms the diagnosis of adenomatous polyposis and allows relatives to be tested with a high degree of accuracy. Once an affected patient has been genotyped, all at-risk relatives can be screened for the mutation.

Surveillance and management of CRC and polyps

Recommendation

8. In individuals at risk for or affected with the classic AP syndromes, screening for CRC by annual colonoscopy or flexible sigmoidoscopy should be performed, beginning at puberty. In families with AFAP or MAP, surveillance should be by colonoscopy (strong recommendation, moderate quality of evidence).

Summary of evidence

Colon screening should be performed in those with a clinical or genetic diagnosis of FAP or in FDRs of those with FAP if genetic testing is uninformative or has not been done. In families where no mutation can be found, all at-risk relatives must undergo endoscopic screening. Colonoscopy should begin at puberty, or whenever there are suggestive

symptoms such as chronic diarrhea, rectal bleeding, or abdominal pain. Flexible sigmoidoscopy is also reasonable in families with classic FAP, until a polyp is found. If this is proved to be an adenoma, full colonoscopy should be done. During colonoscopy, polyp number, size, and distribution should be recorded, and several polyps should be biopsied. The average age of FAP diagnosis in patients presenting with symptoms is 35.8 years (range, 4–72 years) in the St Marks FAP registry in London (81). Polyps begin to appear most often in the second or third decade of life. The mean age of polyp occurrence is 15.9 years (range, 8–34 years) (119). Adenomatous polyps are usually distributed evenly throughout the colon, with a slight distal colonic excess. The size of the polyps depends on the stage at which the patient is examined. Even in fully developed cases, however, 90% of adenomas are <0.5 cm in diameter, and <1% of polyps are >1 cm. Polyps may either carpet the colon with myriad small lesions or occur as more distinct and somewhat larger lesions. Striking heterogeneity of polyp number and growth rate has been observed (120). Histopathology demonstrates tubular adenomas, indistinguishable from common or sporadic adenomas. Villous and tubulovillous histologies are also seen, but much less frequently and usually in larger polyps. A histologic feature of FAP not observed in the general population is dysplastic or adenomatous epithelial cells in single crypts or even portions of single crypts. These are called microadenomas and are often seen in FAP biopsy specimens of normal-appearing mucosa (81). Budding of dysplastic epithelium from normal crypts can be observed and aberrant crypt foci have been reported to occur with increased frequency in FAP (121). These lesions are similar to microadenomas but are identified with methylene blue staining of the colonic mucosal surface.

Colonic adenocarcinoma is the inevitable consequence of FAP unless the colon is removed. There is a 25% incidence of colon cancer in newly diagnosed FAP patients, not belonging to known families, that remains common because of the high frequency of *de novo* germline mutations (94,122). In the St Mark's series, the average age at cancer diagnosis was 39 years. By 45 years of age, 87% had developed cancer, and by 50 years, it increased to 93%. Colon cancer has been reported as early as 9 years of age, although the occurrence of malignancy before adolescence is very unusual. Multiple colonic malignancies were present in ~48% of those with cancer (41% synchronous and 7% metachronous). Of the malignancies, 84% were at or distal to the splenic flexure, a fraction almost identical to that found in their series of random colorectal malignancies at that time. Average life expectancy after diagnosis of cancer was 2.6 years.

The generally accepted colon screening guideline for children at risk for classic FAP is every 1- to 2-year sigmoidoscopy beginning at 10 to 12 years of age (123–127). Those initially screened at an older age should probably have colonoscopy for the first examination. If surgery is delayed longer than a year after polyps emerge, annual colonoscopy should be used for surveillance.

Colon screening with subsequent surgery decreases and almost eliminates mortality from large bowel malignancy in FAP (127–130). Survival is remarkably improved in relatives of probands who undergo screening (82,85,86,129,131).

In AFAP, the emergence of adenomas and cancer is delayed 10 to 20 years compared with typical FAP. The cumulative risk of CRC in AFAP by age 80 years in two large carefully studied kindreds was estimated to be 69%, with an average age at cancer diagnosis of 58 years (range, 29–81 years) (90). In another study, the average age at symptomatic presentation was 52 years (132). For AFAP, colonoscopy should always be used for screening, in view of more proximal colonic polyp distribution. Onset of examination in AFAP can reasonably be delayed until the late teens to mid-20s and be performed every 1 to 2 years.

MAP patients commonly have between 20 and 99 polyps (107,111) and rarely have >500, although the colonic phenotype can vary (107,112,113). CRC was found to be present in patients with an MAP diagnosis in ~60% of cases (107). CRCs in MAP have been predominantly distal colonic in some studies (105) and proximal in others (133). In eight population-based studies, 28 of 79 (35%) MAP cases with CRC had no concurrent polyps, whereas 17 (22%) had <10 adenomas (107). The risk of CRC by age 50 years is 19% and by age 60 years is 43%, with an average age of onset of 48 years (107,134,135). Although the predominant polyp type in patients with MAP is an adenoma, multiple hyperplastic and/or sessile serrated polyps (also referred to as sessile serrated adenomas) may occur. In a small study of 17 patients with MAP, 8 (47%) had at least one hyperplastic and/or sessile serrated polyp, 3 (~18%) met criteria for serrated polyposis (previously referred to as hyperplastic polyposis, see Serrated polyposis section below for additional details), and 1 patient had over 100 hyperplastic and sessile serrated polyps (136). Treatment of MAP follows the same principles as AFAP. The disease may be managed endoscopically with at least yearly colonoscopy. If the polyps become endoscopically uncontrollable, then colectomy is indicated. Currently, there is no consensus as to whether monoallelic *MUTYH* mutations warrant increased CRC screening.

No recommendations for treatment or surveillance of patients with polymerase proofreading-associated polyposis have been made because the frequency of polyps, cancer, and the extracolonic phenotype have yet to be determined, but the options of close endoscopic surveillance and colectomy seem reasonable.

Recommendation

- 9.** Absolute indications for immediate colorectal surgery in FAP, AFAP, and MAP include: documented or suspected cancer or significant symptoms. Relative indications for surgery include the presence of multiple adenomas >6 mm, a significant increase in adenoma number, the presence of an adenoma with high-grade dysplasia, and inability to adequately survey the colon because of multiple diminutive polyps (strong recommendation, low quality of evidence).

Summary of evidence

Development of colon cancer in classic FAP is inevitable if the colon is not removed. An appropriately timed colectomy remains the cornerstone of colon cancer prevention in FAP (127,137,138). Prophylactic surgery can be planned at a suitable time (late teens to early twenties), based on the risk of cancer posed by the polyp burden. Indications for early

surgery include polyps >10 mm diameter, polyps with high-grade dysplasia, marked increases in polyp number from one exam to the next, and symptoms. Surgical options are colectomy with IRA (for <20 rectal and <1,000 colonic adenomas) and proctocolectomy with IPAA (for severe or profuse adenomas, >20 rectal adenomas, and >1,000 colonic adenomas). A laparoscopic approach is now often used for both surgical approaches. Conversion from IRA to IPAA may occasionally be needed because of development of numerous or advanced rectal adenomas. Proctocolectomy with ileostomy is rarely needed. *APC* mutation location, allowing prediction of severity of rectal polyposis and likelihood of future completion proctectomy, has been suggested as a factor to consider in determining which procedure should be done (139).

Colectomy with IRA is a single-stage procedure with slightly less morbidity than the IPAA surgery, but some rectal cancer risk remains and yearly proctoscopy is essential (139,140). Even after total proctocolectomy and IPAA, adenomas and cancers may occur in the anal transition zone and in the pouch itself; lifelong endoscopic surveillance is required (141–143). Possible morbidities from either surgery include increased bowel frequency and incontinence. Pouch surgery is associated with some loss of fertility in women and some loss of sexual function in men.

Patients with AFAP can often be managed for many years with colonoscopic polypectomy and may possibly never need colectomy (90). If surgical resection is indicated, AFAP patients can almost always undergo colectomy and IRA because of rectal sparing of polyps. After colectomy with IRA in a large series of patients with AFAP, an average of 3.4 recurrent polyps (range, 0–29) and only one cancer was found in the postcolectomy rectal remnant over a mean follow-up of 7.8 years (range, 1–34 years) (90).

Similar to AFAP, FAP, and some multiple adenoma patients, subtotal colectomy with close subsequent surveillance would seem to be the best option for MAP patients with relative rectal sparing. Restorative proctocolectomy is indicated if the rectum is substantially involved.

Surveillance and management of extracolonic malignancies

Recommendations

10. Screening for gastric and proximal small bowel tumors should be done using upper endoscopy including duodenoscopy starting at age 25–30 years. Surveillance should be repeated every 0.5–4 years depending on Spigelman stage of duodenal polyposis: 0=4 years; I=2–3 years, II=1–3 years, III=6–12 months, and IV=surgical evaluation. Examination of the stomach should include random sampling of fundic gland polyps. Low-grade dysplasia is common in fundic gland polyps, and surgery should be reserved for high-grade dysplasia or cancer (strong recommendation, very low quality of evidence).
11. Annual thyroid screening by ultrasound should be recommended to individuals affected with FAP, MAP, and attenuated polyposis (conditional recommendation, low quality of evidence).

12. Biannual screening should be offered to affected infants annually until age 7 years with α -fetoprotein and ultrasounds (conditional recommendation, very low quality of evidence).

Summary of evidence

The phenotype of FAP includes benign and malignant neoplasms in other organs. Other organs commonly affected include the thyroid (with papillary thyroid cancer), adrenal (non functioning adenomas), the small intestine (adenomas or carcinoma), bones (osteomas), retina (congenital hypertrophy of the retinal pigmented epithelium), and skin (epidermoid cysts). However, the most common causes of death in FAP after CRC are duodenal or ampullary cancer and desmoid disease.

Endoscopically visible duodenal adenomas are found in more than half of FAP patients (101,144,145). The lifetime risk for duodenal cancer is 3 to 5%, but in some series it has been even higher (81,101,127,144,146). The age of duodenal cancer diagnosis ranges from 17 to 81 years, with a mean between 45 and 52 years. Approximately half of duodenal cancers are ampullary or periampullary, whereas others are elsewhere in the duodenum (147). Duodenal cancer is one of the leading causes of death in FAP patients who have had prophylactic colectomy (147–150).

Adenomas beyond the duodenum may occur throughout the small bowel but are concentrated for the most part in the proximal jejunum (50% of cases) and distal ileum (20% of cases) (151–153). The polyps are most commonly 1 to 10 mm in diameter and multiple. Most duodenal polyps cluster around the ampulla, although in some patients there are small adenomas scattered throughout the duodenum. Adenomas may progress, often slowly, and there is evidence of an adenoma–carcinoma sequence similar to that observed in the colon (101). Adenomas sometimes grow large, exhibit villous histology and increasing degrees of dysplasia, and may cause symptoms. A scoring system has been developed to evaluate the severity of duodenal polyposis and is now widely applied as the Spigelman staging system (154) (**Table 9**). The risk for duodenal cancer increases to 36% within 10 years for Spigelman stage IV patients (155). The risk of exhibiting Spigelman stage IV duodenal polyposis is 43% by age 60 years and 50% by age 70 years (156). Patients with Spigelman Stage IV duodenal adenomatosis are candidates for a pancreas-preserving duodenectomy. This is much less morbid than a Whipple procedure and patients have a better quality of life. If there is a strong suspicion of cancer, then a Whipple is necessary.

Gastric fundic gland polyps are also common but gastric adenomas are rare and in western countries gastric cancer is uncommon. Gastric polyps occur in 23 to 100% of FAP patients (101,126,144,157). In the gastric fundus and body, the polyps are most often fundic gland polyps, considered hamartomas. These polyps are histologically seen to consist of simple hyperplasia of the fundic glands with microcysts. Endoscopically, the polyps are multiple sessile lesions, most often 1 to 10 mm in diameter, and are the same color as surrounding mucosa (158). Considerable variation in size and number is observed. The polyps are sometimes so numerous that they coalesce, forming areas of irregular, matted surface mucosa. Fundic gland polyps rarely cause symptoms. Almost half of FAP patients with fundic gland polyps will have superficial dysplasia in some of those polyps (159). Although

they are considered nonneoplastic, fundic gland polyps may rarely progress to cancer (160–162). Adenomatous polyps occur in the stomach of ~10% of patients with FAP. They are most often confined to the antrum but are occasionally found in the body and fundus. (163,164) The lifetime risk for gastric cancer in FAP is ~0.6%, believed both from fundic gland polyps and adenomatous polyps (138).

Upper GI screening has not been demonstrated to improve prognosis but is nonetheless recommended in view of the cancer risk and expectation that mortality can be improved (127,144–147,156,165). Standard upper endoscopy should be supplemented with a side-viewing instrument to visualize the duodenal papilla. Duodenal screening should begin at age 25–30 years and continue for life, with a frequency determined by the severity of the duodenal polyposis as measured by the Spigelman score. A 0.5–4-year interval for examination is given as follows: (i) every 4 years for Spigelman stage 0; (ii) every 2–3 years for stage I disease; (iii) every 1–3 years for stage II disease; (iv) every 6–12 months for stage III disease; and (v) for stage IV disease: surgical evaluation, expert surveillance every 3–6 months and complete mucosectomy or duodenectomy, or Whipple procedure if duodenal papilla is involved (101,127). Another approach to screening is every 3-year endoscopy if adenomas are not found and annually if they are. The stomach should be examined during endoscopy and any polyps judged to be of concern because of size, color, or gross appearance biopsied. The role of examination of the small bowel beyond reach of the upper endoscope by computed tomography (CT) enterography, push or balloon enteroscopy, or capsule endoscopy if upper endoscopy demonstrates severe duodenal polyposis is uncertain (153).

In contrast to colorectal polyps and cancer, the expression of upper GI polyps, both gastric and duodenal, does not appear to be attenuated in number, age at emergence, or cancer risk in AFAP compared with FAP (89,162); therefore, EGD surveillance should also be performed at age 25–30 years, and continued according to the rules stated for classical FAP.

Given that the risk of duodenal cancer in MAP is similar to that of AFAP and FAP, upper GI endoscopy with added side-viewing duodenoscopy should be considered, starting at around age 30 years and repeated at intervals similar to AFAP and FAP, again depending on duodenal findings (166).

Gallbladder, bile ducts, and pancreas

Both adenomatous change and cancer have been reported in the gallbladder, bile ducts, and pancreas (138,151,167–170). Biliary and pancreatic duct obstructions have arisen from both benign and malignant lesions. The cancer risks are shown in **Table 7**. There are no surveillance strategies that are currently recommended for these malignancies.

Extraintestinal malignancies

Up to 12% of FAP patients have thyroid cancer and 80% have nodular thyroid (171). The mean age of diagnosis of thyroid cancer is 28 years, ranging from 12 to 62 years (172). A female preponderance is observed, and the histology is predominantly papillary, commonly

with a cribriform–morular pattern. Annual thyroid ultrasound is recommended for thyroid screening in FAP (171).

Hepatoblastoma occurs in 1.6% of FAP patients, exhibits a male predominance, and associates somewhat with mutations in the 5' end of the *APC* gene. This malignancy most often occurs in the first 5 years of life, with some risk up to 15 years of age (173). Screening with every 3- to 6-month serum α -fetoprotein and liver ultrasound for the first 5 to 10 years in FAP patients has been suggested but is still debated. A family history of hepatoblastoma may be an indication to do this from age 6 months to 6 years. A decision to perform hepatoblastoma screening mandates genetic testing in infancy to see if the child carries the mutation.

Desmoid tumors

Screening is not done for desmoids, but evaluation is done for palpable masses and a full work-up for suggestive symptoms. Periodic abdominal imaging is not generally recommended, but preoperative abdominal CT scan before colectomy may be considered if desmoids have been an issue in family members.

The lifetime risk of extracolonic tumors in MAP is not as well defined as the colorectal phenotype. In a large study of 276 MAP patients, 17% had extracolonic lesions, with an estimated 38% lifetime risk of extracolonic malignancy that is approximately double the risk in the general population (174). Similar to FAP and AFAP, the lifetime risk of duodenal cancer in MAP has been estimated to be 4% (174). Although gastric lesions have been found in up to 11% of patients with MAP, data are currently lacking to support an increased risk of gastric cancer (107).

Other cancers such as endometrial, breast, ovarian, bladder, various skin, and thyroid have been reported in patients with MAP (107), although it is still not clear whether the lifetime risk for these malignancies is increased. Although rare, other findings seen in patients with MAP have included sebaceous gland adenomas, carcinomas and epitheliomas, lipomas, congenital hypertrophy of the retinal pigment epithelium, osteomas, desmoid tumors, epidermoid cysts, and pilomatrixomas (107). Surveillance and disease management of the colon in MAP should be similar to patients with multiple adenomas, AFAP, and FAP (166).

In women with a *POLD1* mutation, pelvic ultrasound and selective endometrial biopsy may be considered as the *POLD1* variant has been associated with endometrial and possibly brain tumors.

Recommendation

13. Postsurgical surveillance should include yearly endoscopy of rectum or ileal pouch, and examination of an ileostomy every 2 years (strong recommendation, low quality level of evidence).

Summary of evidence

Adenomas may develop in the ileal pouch after colectomy with IPAA surgery, or they may develop in the small segment of remaining rectal epithelium after restorative

proctocolectomy (175–177). There appears to be a small but real risk for cancer in the ileal pouch (177–179). Advanced dysplasia and cancer may occur at the anal transition zone, either from rectal tissue unexpectedly remaining with ileo-anal anastomosis or from the short segment of rectum often remaining with restorative proctocolectomy (143,150,180).

After colectomy or proctocolectomy, endoscopic surveillance of the rectum or ileal pouch should continue yearly. Concerning polyposis in the rectum (large adenomas, high-grade dysplasia, >20 adenomas) is treated either by polypectomy or proctectomy. Pouch polyposis can be treated by polypectomy or chemoprevention with sulindac. Ileostomies should be checked every 2 years as adenomas and even cancer can develop on the stoma.

Prevention strategies

Much attention and effort has been given to examining chemoprevention for colonic and duodenal polyps in FAP (127). Considerable regression and prevention of colonic and rectal adenomas has been demonstrated with sulindac, but cancer prevention is less certain. Celecoxib appears to have a more modest effect in the colon and rectum, but some effect in duodenal adenoma regression as well. Celecoxib was approved for use in the United States for several years for FAP, but this indication has now been removed. Concern over cardiovascular side effects of long-term cyclooxygenase-2 (COX-2) inhibitors has dampened enthusiasm for their use in FAP. In view of the uncertainty of cancer prevention with sulindac, it is not considered a substitute for colectomy but has shown utility in rectal surveillance by substantially decreasing the number of adenomas needing removal at periodic examination. Chemoprevention studies examining nonsteroidal anti-inflammatory drugs and other agents continue in the hopes that colectomy might be delayed.

Hamartomatous polyposis syndromes

Peutz–Jeghers syndrome

Indications for genetic testing: Summary statement

- Individuals with perioral or buccal pigmentation and/or two or more histologically characteristic GI hamartomatous polyp(s) or a family history of PJS should be evaluated for PJS.

Summary of evidence

PJS is an autosomal-dominantly inherited syndrome that includes histologically distinctive hamartomatous polyps of the GI tract and characteristic mucocutaneous pigmentation (181–184). Its incidence is estimated at between 1 in 50,000 and 1 in 200,000 births (185). The mucocutaneous melanin pigment spots are seen in >95% of cases. They are 1 to 5 mm in diameter and most commonly occur in the perioral area and on the buccal mucosa (94%). Pigment spots on the lips are distinctive in that they cross the vermilion border and are often much darker and more densely clustered than common freckles. These spots also occur on the face, forearms, digits, palms, soles, perianal area, and rarely on the intestinal mucosa. The pigment appears in infancy and may fade with age, but less so on the buccal mucosa. GI polyps occur in 88 to 100% of patients. PJS polyps are histologically distinct. They are nondysplastic, have normal overlying epithelium specific to the GI segment in which they

are found, and exhibit an arborizing pattern of growth with muscularis mucosae extending into branching fronds of the polyp. Epithelial infolding may result in what is termed *pseudoinvasion* that can lead to an incorrect diagnosis of cancer. Adenoma and cancer may occur in PJS polyps (186). Their frequency by segment is: stomach, 24%; small bowel, 96%; colon, 27%; and rectum, 24% (183,186). Polyp sizes range from 0.1 to 3 cm in diameter. Polyp growth begins in the first decade of life, but patients typically do not develop symptoms until the second or third decade (187,188). Symptoms arise from larger polyps that may infarct, ulcerate, bleed, and cause intestinal obstruction and intussusception, usually in the small intestine.

Genetic etiology

Summary statement

- Genetic evaluation of a patient with possible PJS should include testing for *STK11* mutations.

Summary of evidence

PJS arises from mutations of the *STK11* gene, a tumor suppressor seronine/threonine kinase gene, previously called *LKB1*, on chromosome 19p (182,183). Up to 94% of PJS families have mutations of *STK11* with up to a third of disease causing mutations representing large deletions (189,190). Approximately 25% of newly diagnosed PJS patients represent *de novo* mutations (183). There do not appear to be genotype–phenotype correlations with mutation location in the *STK11* gene (191). Once a disease-causing mutation is identified in a patient with PJS, other family members should undergo mutation-specific testing to determine whether the disease is present or absent so that appropriate surveillance can be undertaken.

Surveillance and management

Recommendation

14. Surveillance in affected or at-risk PJS patients should include monitoring for colon, stomach, small bowel, pancreas, breast, ovary, uterus, cervix, and testes cancers. Risk for lung cancer is increased, but no specific screening has been recommended. It would seem wise to consider annual chest radiograph or chest CT in smokers (conditional recommendation, low quality of evidence).

Summary of evidence

Numerous studies and reviews have now reported a high risk of both GI and extraintestinal cancer in PJS (126,182,183,185,192–197). Individual risks by cancer site are given in **Tables 5** and **7**. The malignant risk in PJS includes colorectal, breast, pancreatic, gynecological, small bowel, lung, and gastroesophageal cancers in that order of risk (198). The overall risk of developing any cancer at ages 20, 30, 40, 50, 60, and 70 years was 1%, 3%, 19%, 32%, 63%, and 81% respectively. In terms of specific cancers, estimated lifetime risks are 39% for colorectal, 29% for gastric, 13% for small bowel 24–54% for breast, 21% for ovary, 10–23% for cervix, 9% for uterus, 9% for testicular, 7–17% for lung, and 11–36% for pancreas (196). Distinctive tumors in women with this condition include ovarian sex

cord tumors with annular tubules that are benign, although ~20% become malignant; mucinous tumors of the ovary; and well-differentiated adenocarcinomas of the uterine cervix, called adenoma malignum (186). Nine percent of males develop large cell calcifying sertoli cell tumors of the testes, resembling sex cord tumors with annular tubules, that have a 10 to 20% chance of becoming malignant (199). Feminization may occur with the benign testicular tumors. Mainly because of cancer, the overall survival of PJS patients is significantly shorter than age- and gender-matched controls (200).

Surveillance guidelines for PJS are empiric and based on the risk for GI complications and cancer. See **Table 10** for specific recommendations. A consortium review group has recommended that upper GI endoscopy (EGD) and colonoscopy be done first at age 8 years (182). If polyps are found, both examinations should be repeated every 3 years. If none are found, a second baseline examination should be done at age 18 years and then every 3 years there-after. Similar surveillance is recommended for the small bowel, i.e., first examine the small bowel by video capsule endoscopy at age 8 years, but then repeat this surveillance every 3 years from that age. Modern CT enterography is accurate at detecting small bowel polyps, particularly those ≥ 1 cm in diameter, but repeated X-ray exposure is problematic.

Treatment involves EGD and colonoscopic removal of polyps (probably all those >0.5 or 1 cm in diameter) (185). Clearing of all polyps is preferable but not always possible. Colectomy is sometimes necessary to control colonic polyps and should be considered if colonoscopic management is difficult and especially if neoplastic change is found in colonic polyps. Intussusception is the primary complication of small bowel polyps, starting at a young age, and continuing throughout life (188). Surveillance and treatment of the small bowel are based in large part on prevention of this complication. In the recent study by van Lier *et al.* (188), the initial episode of intussusception occurred at a median age of 16 years (range, 3–50 years), with 50% of first episodes presenting by age 20 years. Of all intussusceptions, 80% presented as an acute abdomen and the average polyp size causing this complication was 3.5 cm (range, 1.5–6.0 cm). When small bowel intussusceptions occur, surgery is often necessary and should include careful examination of the entire small bowel to eliminate all significant polyps. Intraoperative endoscopy is often a helpful adjunct to accomplish extensive polyp removal. This is also an appropriate time to examine and remove gastric and duodenal polyps of significant size (201). The advent of video capsule endoscopy, double balloon enteroscopy, and CT enterography is changing diagnostic and management approaches to PJS by allowing earlier detection of polyps and non-operative removal in many cases (202–205).

Prevention strategies

Chemoprevention approaches to decrease polyp burden in PJS are under study but not yet a reality. PJS polyps exhibit overexpression of COX-2, suggesting that COX-2 inhibitors may be useful in reducing polyps (206). Hyperactivation of the mammalian target of rapamycin has been associated with PJS. In addition, inhibition of mammalian target of rapamycin in a PJS mouse model has demonstrated decreased polyp burden (207). Everolimus, a mammalian target of rapamycin inhibitor, is under study as a potential agent for treatment of PJS (182).

Juvenile polyposis syndrome

Indications for genetic testing—*Summary statement*

- Individuals with five or more juvenile polyps in the colorectum or any juvenile polyps in other parts of the GI tract should undergo evaluation for JPS.

Summary of evidence

JPS is an autosomal-dominantly inherited condition where multiple juvenile polyps are found in the colorectum (98%), stomach (14%), jejunum and ileum (7%), and duodenum (7%) (183,208–210). The incidence of JPS is between 1 in 100,000 and 1 in 160,000 individuals (210). The polyps in JPS vary in size from small sessile nodules to pedunculated lesions that are 3 cm in diameter. Most large polyps are pedunculated, but small polyps, especially those in the stomach, are sessile. Grossly, most polyps exhibit a surface that is smooth, rounded, reddish colored, and without fissures or lobulations; large polyps may appear to be multilobulated. A white exudate is often seen on the polyp surface. On cut section, there are cystic spaces filled with mucin. Microscopically, there is abundant lamina propria with benign but often elongated and cystically dilated glands and lack of a smooth muscle core. Excess chronic inflammatory cells are sometimes present. The epithelial lining of the surface and cysts is nondysplastic and reflects the area of the GI tract where the polyp is located. Polyps begin to appear in the first decade of life, and dozens to many hundreds of polyps are present in the fully developed syndrome. Most patients develop symptoms in the first two decades of life. The average age at diagnosis is 18.5 years but may be later. Rectal bleeding with anemia is the most common presenting symptom, followed by abdominal pain, diarrhea, passage of tissue per rectum, and intussusception (183,208). The majority of colonic polyps, 70% in one study, occurred in the proximal colon (210).

The generally accepted clinical criteria for JPS include: (i) at least five juvenile polyps in the colorectum; (ii) juvenile polyps in other parts of the GI tract; or (iii) any number of juvenile polyps in a person with a known family history of juvenile polyps (210).

Genetic etiology

Summary statement

- Genetic evaluation of a patient with possible JPS should include testing for *SMAD4* and *BMPRIA* mutations.

Summary of evidence

Juvenile polyposis occurs as a result of mutations of the *SMAD4* gene (also called the *MADH4* gene) or the *BMPRIA* gene (210–214). Up to 60% of individuals with clinically defined JPS are now found to exhibit mutations of the *SMAD4* or *BMPRIA* genes (215). Approximately 25% of newly diagnosed cases are sporadic and thus represent new or *de novo* mutations, whereas 75% will have a family history (183). Fourteen percent of mutations are large deletions and 10% are promoter mutations (215–217). Both genes are tumor suppressor genes involved in the tumor growth factor- β signaling family and directly or indirectly affect cell growth inhibition and apoptosis. There is evidence that normal bone morphogenetic protein signaling also suppresses Wnt signaling to ensure a balanced control

of stem cell self-renewal (218). Biallelic gene inactivation has been noted in both stromal cells and epithelial cells of polyps (208,219).

Genetic testing is particularly important in JPS, both to confirm the diagnosis in a proband and to test relatives. Testing is also important to separate JPS from other conditions in which juvenile polyps form, especially CS and Bannayan–Riley–Ruvalcaba syndrome. Once a disease-causing mutation is identified in a patient with JPS, other family members should undergo mutation-specific testing to determine whether the disease is present or absent so that appropriate surveillance can be undertaken.

Surveillance and management

Recommendations

15. Surveillance of the GI tract in affected or at-risk JPS patients should include screening for colon, stomach, and small bowel cancers (strong recommendation, very low quality of evidence).
16. Colectomy and IRA or proctocolectomy and IPAA is indicated for polyp-related symptoms, or when the polyps cannot be managed endoscopically (strong recommendation, low quality of evidence).
17. Cardiovascular examination for and evaluation for hereditary hemorrhagic telangiectasia should be considered for *SMAD4* mutation carriers (conditional recommendation, very low quality of evidence).

Summary of evidence

JPS mutation carriers have a very high risk for colon cancer and an increased risk for gastric, duodenal, and pancreatic cancers (**Tables 5 and 7**). The cancer risk in JPS is believed to arise from adenomatous tissue within the juvenile polyp, as up to 50% of juvenile polyps in JPS contain areas of adenomatous change. The risk of colon cancer is 17–22% by age 35 years and approaches 68% by age 60 years (183,220). The mean age of colon cancer is 34 years, with a range of 15 to 68 years. Gastric cancer risk is 30% in those with *SMAD4* mutations (183,210). The median age of upper GI carcinoma is 58 years, with a range of 21 to 73 years (221,222).

Surveillance guidelines for JPS are found in **Table 10**. Colonoscopy should be annual, beginning at age 12 years or earlier if symptoms occur, especially rectal bleeding. It should be repeated every 1 to 3 years depending on polyp burden and polyps ≥ 5 mm should be removed (210). Upper endoscopy is recommended every 1 to 3 years beginning at age 12 years, or earlier for symptoms, and should be repeated every 1 to 3 years, depending on severity with removal of polyps ≥ 5 mm. The small bowel past the duodenum should be periodically surveilled, depending on initial polyp findings, by enteroscopy, capsule endoscopy, and/or CT enterography if duodenal polyposis is present or if there is unexplained anemia, protein-losing enteropathy, or other small bowel symptoms. Patients with limited numbers of polyps in any area of the GI tract can usually be managed with endoscopic polypectomy. Colectomy with IRA is indicated if cancer, high-grade dysplasia, or polyposis cannot be adequately controlled endoscopically (183,208). Surveillance of the

remaining rectum or pouch is necessary (223). Proctocolectomy with IPAA may be needed depending on the number of rectal polyps (223). Half of those with IRA will later need proctectomy because of polyp perfusion. Complete or partial gastrectomy may also be necessary for patients with advanced dysplasia, gastric cancer, or even massive gastric polyps that cannot be effectively controlled endoscopically (183,208). Other screening should include annual complete blood count, cardiovascular examination, and hereditary hemorrhagic telangiectasia protocol evaluation if *SMAD4* mutation is present (210).

Cowden syndrome (PTEN hamartoma tumor syndrome)

Summary statement

- Individuals with multiple GI hamartomas or ganglioneuromas should be evaluated for CS and related conditions.

Summary of evidence

CS and its variants, including Bannayan–Riley–Ruvalcaba syndrome and PTEN hamartoma tumor syndrome (PHTS), have been associated with a broad range of clinical phenotypes. Colonic polyps are found in up to 95% of CS patients undergoing colonoscopy (224,225). Polyps are few to numerous (even hundreds) and are distributed throughout the colon. The natural history of polyps is not well characterized, although polyps may occur at a young age. Hamartomatous polyps are the most common histologic type, occurring in up to 29% in one study (224). Polyp types include juvenile polyps, ganglioneuromas, adenomas, and inflammatory polyps (224,226,227), and less commonly leiomyomas, lipomas, and lymphoid polyps (228). Hyperplastic polyps have also been reported as an association, but have not been observed in all studies (224,227). The majority of CS patients have multiple synchronous histologic types at colonoscopy.

A frequent finding in the esophagus is diffuse glycogenic acanthosis (228,229). One or several such lesions may occasionally be observed in patients undergoing EGD for various reasons, but diffuse, sometimes many hundreds of lesions are observed in 80% of those with PHTS (227). It has been suggested that diffuse esophageal glycogenic acanthosis combined with colonic polyposis should be considered pathognomonic for CS (228).

Several investigations report the frequent finding of multiple hamartomatous polyps in the stomach, duodenum, and small bowel (183,224,229). Similar to the colon, histologies include hamartomas, hyperplastic polyps (different from colonic hyperplastic polyps), ganglioneuromas, adenomas, and inflammatory polyps. An upper GI study of 10 *phosphate and tensin homolog (PTEN)* mutation-positive patients found all 10 to have multiple hyperplastic gastric polyps and 3 to have multiple hamartomatous polyps in that location (227). One patient had a single hamartomatous polyp in the duodenum whereas three had adenomatous polyps. There are reports of gastric and colon cancers in CS patients (183,230).

Specific indications for evaluation for CS are delineated in **Table 11**.

Genetic etiology

Summary statement

- Genetic evaluation of a patient with possible CS should include testing for *PTEN* mutations.

Summary of evidence

CS is caused by mutations in the *PTEN* gene (231–234). Once a disease-causing mutation is identified in a patient with CS or related conditions, other family members should undergo mutation-specific testing to determine whether the disease is present or absent so that appropriate surveillance can be undertaken.

Surveillance and management

Recommendation

18. Surveillance in affected or at-risk CS patients should include screening for colon, stomach, small bowel, thyroid, breast, uterine, kidney, and skin (melanoma) cancers (conditional recommendation, low quality of evidence).

Summary of evidence

Management of PHTS involves prevention and early detection of the associated cancers through surveillance. Colon cancer has not been associated with CS historically (183), although recent studies have indeed shown increased risk for this malignancy. One multicenter study found 13% of *PTEN* mutation carriers to have colon cancer, all younger than 50 years of age (224). Investigations have now indicated a 9 to 16% lifetime risk for large bowel cancer (225,235,236). It is uncertain whether colon malignancy arises from adenomatous and/or hamartomatous polyps in PHTS, although there is little doubt as to the increased risk and possibility of young age onset.

Surveillance recommendations are provided in **Table 10**. Recommendations are all expert opinion based rather than evidence based, and derived from screening guidelines of the relevant cancers in other settings but adjusted for the malignancy risks observed in PHTS.

Hereditary mixed polyposis syndrome (HMPS) is a condition that was originally described in a large Ashkenazi Jewish family with multiple colorectal polyps and cancer. Affected patients exhibited mixed juvenile–adenomatous polyps and also adenomatous, hyperplastic, serrated adenomas, and mixed hyperplastic–adenomatous polyps and adenocarcinomas. Mean age of polyp occurrence in one family was 28 years. HMPS may be misdiagnosed as JPS or serrated polyposis syndrome (SPS) and vice versa.

Even though HMPS linked to a locus on chromosome 15q13.3–q14 in a number of families, which includes the *CRAC1* gene, the etiology remains elusive. Recently, a duplication 40 kb upstream of the *GREM1* gene locus at chromosome 15 was found in two individuals with HMPS. The authors hypothesized that this duplication interacts with the *GREM1* promoter causing increased *GREM1* expression, resulting in a predisposition to multiple colorectal polyps.

Genetic testing for *GREM1* mutation and expression might be considered in families with adenomatous and hamartomatous polyposis in which an etiology cannot be determined. Management from what is now known should probably be similar to that for FAP, depending on the polyp number, size, and histology.

Serrated polyposis syndrome

Clinical definitions—*Summary statement*

- Individuals who meet at least one of the following criteria have the clinical diagnosis of SPS: (i) at least 5 serrated polyps proximal to the sigmoid colon with 2 of these being >10 mm; (ii) any number of serrated polyps proximal to the sigmoid colon in an individual who has a first-degree relative with serrated polyposis; and (iii) >20 serrated polyps of any size, distributed throughout the large intestine.

Summary of evidence

SPS, previously referred to as hyperplastic polyposis syndrome, is a rare condition currently defined by clinical criteria and characterized by a predisposition to serrated polyps and an increased risk of CRC (237–239). Originally, hyperplastic polyps were the only lesion included in the diagnostic criteria for hyperplastic polyposis (240). In addition to hyperplastic polyps, other serrated polyps including sessile serrated polyps and traditional serrated adenomas may also be found, and hence the preferred term serrated polyposis syndrome (241).

The updated World Health Organization (WHO) diagnostic criteria for SPS include any one of the following: (i) at least five serrated polyps proximal to the sigmoid colon with two or more of them >10 mm in diameter, (ii) any number of serrated polyps proximal to the sigmoid colon in an individual who has a FDR with SPS, or (iii) >20 serrated polyps of any size, but distributed throughout the colon (242). The true prevalence of SPS is not known, but had been previously estimated to be 1:100,000 based on a large screening colonoscopy study of 50,148 participants in which 28 subjects (0.06%) were found to have the syndrome. (243). More recent studies have evaluated the prevalence of SPS based on the WHO criteria. The National Health Service Bowel Cancer Screening Programme (NHSBCSP) reviewed all pathology and colonoscopy records for guaiac fecal occult blood test-positive patients presenting for index screening colonoscopy (244). Out of 755 patients, 5 (0.66%) met SPS criteria 1 and/or 3 (244). There-fore, 1 in 151 patients in the NHSBCSP met SPS criteria during their index colonoscopy, a much higher rate than previous reports of SPS in the general population (244). In a study from Barcelona, the prevalence of SPS in patients undergoing colonoscopy after a positive fecal immunochemical testing was 8 out of 2,355 (0.34%) (245).

Genetic etiology

Summary statement

- Indications for genetic testing. A clear genetic etiology has not yet been defined for SPS, and therefore genetic testing is currently not routinely recommended for SPS

patients; testing for *MUTYH* mutations may be considered for SPS patients with concurrent adenomas and/or a family history of adenomas.

Summary of evidence

Although the genetic etiology of SPS remains unknown, 3 patients were found to meet diagnostic criteria for SPS in a series of 17 biallelic *MUTYH* mutation carriers (18%) (136). Conversely, only one biallelic *MUTYH* mutation carrier was observed in a study of 126 patients with SPS (0.8%) (246). In both studies, the patients who met criteria for SPS also reported a history of adenomas. These observations indicate some overlap in the presentation (and potentially the pathogenesis) of MAP and serrated polyposis. Although data are currently limited, it may be reasonable to consider genetic testing for *MUTYH* mutations in patients with SPS, particularly if adenomas are concurrently seen.

Although the mechanisms are not entirely clear, there seems to be a strong association between smoking and SPS (247). In a small study of 32 SPS patients, the rate of current smoking was 47%, and this was significantly higher than the rate in colonoscopy controls (17%) and population controls (12%) (247). In another study of patients with multiple serrated polyps, many of whom met criteria for SPS, 51 of 88 (58%) were ever smokers (248). It is speculated that smoking, although not the cause of SPS, does intensify the phenotypic expression and therefore may be a modifiable risk factor for colorectal lesions(247).

Various studies have shown that a family history of colorectal and other cancers is common and even increased in patients with SPS. Some have suggested that this supports a hereditary etiology to SPS (249,250). However, nongenetic causes, referral bias, and chance occurrence should not be overlooked as substantial factors in these studies (251).

Surveillance and management of CRC

Recommendation

19. Patients with serrated polyposis should undergo colonoscopies every 1–3 years with attempted removal of all polyps >5 mm diameter (conditional recommendation, low quality of evidence).

Summary of evidence

It is now well established that SPS is associated with an increased risk for CRC (252). The specific lifetime risk of colorectal in SPS is not well defined as most study cohorts to date are relatively small, phenotypically diverse, and subject to referral bias. It has been estimated that the lifetime CRC risk is >50%, although this is likely an overestimate (243). CRC was diagnosed in 5 of 77 (6.5%) SPS patients after a median follow-up time of 1.3 years (253). Four out of five of these CRCs were found in serrated polyps <20 mm (253). Boparai *et al.* (253) estimated that the 5-year risk of CRC under surveillance was 7%. In two large descriptive studies of SPS patients, the majority of index cases displayed a pancolonic distribution of polyps (89–96%), the presence of adenomas (78–80%), a diagnosis of CRC (31–42%), and a mean age at diagnosis of 48 years (250,254). In a smaller prospective series of 13 hyperplastic polyposis syndrome patients, 5 (71%) of the 7 CRCs reported were

located in the right colon (237). In another retrospective study of 77 patients with SPS, 22 (28.6%) were diagnosed with CRC at their baseline colonoscopy. Five (6.5%) more patients developed CRC while being followed for hyperplastic polyposis syndrome (253). Synchronous adenomas are found in the majority of patients with SPS (248,252). Conventional adenomas were more frequently found in patients with CRC than those without (254). In a study of CRCs from SPS patients, 18 of 39 (46%) had the *BRAF* V600E mutation, 2 of 40 (5.0%) had *KRAS* mutations, and 17 of 45 (38%) had loss of immunohistochemistry expression of *MLH1* and *PMS2* (255).

There are no available studies regarding the effectiveness of surveillance in SPS. Based on the reported colon cancer risks, colonoscopy and polypectomy is recommended for individuals who fulfill the WHO definition of SPS. Complete clearance of all polyps ≥ 1 cm should be done when possible. Subsequent colonoscopy intervals should be determined by the number and size of polyps, as well as the number of concurrent adenomas, but generally should be performed every 1–3 years.

Recommendation

20. Indications for surgery for SPS include an inability to control the growth of serrated polyps, or the development of cancer. Colectomy and IRA is a reasonable option given the risks of metachronous neoplasia (conditional recommendation, low quality of evidence).

Summary of evidence

Surgery is advised when polyps cannot be endoscopically controlled (256). Prophylactic or therapeutic colectomy is indicated with an inability to control the growth of serrated polyps, the presence of high-grade dysplasia in a serrated polyp that cannot be removed in its entirety, or the development of cancer. Subtotal colectomy and IRA is a reasonable option given the risks of metachronous neoplasia.

Surveillance and management of extracolonic malignancies

Recommendation

21. There is no evidence to support extracolonic cancer surveillance for SPS at this time. Screening recommendations for family members are currently unclear pending further data and should be individualized based on results of baseline evaluations in family members (conditional recommendation, very low quality of evidence).

Summary of evidence

The data on extracolonic cancers in SPS are insufficient at this time, although Win *et al.* (250) reported an increased risk of colorectal and pancreas cancer in relatives of patients with SPS. However, PC was not found in any of the 115 patients with SPS in a study by Kalady *et al.* (249). In a larger study of 105 patients with SPS and 341 FDRs, no increased risk of extracolonic malignancies were seen in patients or their relatives (257). Hazewinkel

et al. (257) concluded that extracolonic cancer in SPS and their FDRs is not increased compared with the general population.

Familial cases of SPS have been reported, although infrequently in most studies. However, a recent prospective study of 78 FDRs of patients with SPS found that the incidence of SPS was 32% (258). Only one of these relatives was diagnosed with CRC during screening colonoscopy (258). Other studies have reported a family history of CRCs in 50–59% of FDRs of patients with SPS (241,248). In one of these studies, only 2 (5%) of the 38 probands with SPS reported a family history of SPS (241). In a recent study, Boparai *et al.* (259) FDRs of serrated polyposis patients had five times the incidence of CRC, suggestive of a hereditary disorder.

Surveillance recommendations for individuals with a family history of serrated polyposis are currently unclear, pending further data. It is reasonable to screen FDRs at the youngest age of onset of serrated polyposis diagnosis (after the exclusion of other genetic causes), and subsequently per colonoscopic findings (248). The frequency and age of initiation of colonoscopic screening in at-risk family members of patients with SPS is less clear. Oquinea *et al.* (258) recommended that colonoscopy in FDRs start at age 35 years (or 10 years before the youngest age of SPS onset in the family, whichever comes first) and then every 3–5 years if no polyps are found. If the FDRs are found to meet SPS criteria 1 or 3, annual surveillance should be performed or every 2 years if only SPS criteria 2 is met (258). On the contrary, the National Comprehensive Cancer Network (NCCN) recommends that FDRs should have colonoscopy at the earliest of the following: (i) age 40 years, (ii) same age as the youngest SPS diagnosis in the family, and (iii) 10 years before CRC in the family in a patient with SPS (260). Further work is ongoing to better define the cancer risks in probands and their relatives so that accurate risk stratification and risk recommendations can be made regarding SPS.

Prevention strategies

Summary of evidence—Observational studies suggest a strong association between smoking and SPS. Independent studies have reported significant smoking history in patients with SPS, leading to the assumption that smoking may be a modifiable risk factor in the pathogenic pathway of colorectal lesions (248,250).

Hereditary pancreatic cancer

Clinical definitions—*Summary statement*

- Individuals should be considered to be at risk for familial pancreatic adenocarcinoma if they: (i) have a known genetic syndrome associated with pancreatic cancer, including hereditary breast–ovarian cancer syndrome, familial atypical multiple melanoma and mole syndrome (FAMMM), PJS, LS, or other gene mutations associated with an increased risk of pancreatic adenocarcinoma; (ii) have two relatives with pancreatic adenocarcinoma, where one is a FDR; (iii) have three or more relatives with pancreatic cancer; or (iv) have a history of hereditary pancreatitis.

Summary of evidence

The known hereditary syndromes associated with an increased risk for developing PC along with their relative risk for developing PC are shown in **Table 12**. PC is one of the key cancers used in determining whether a patient's family history warrants genetic risk evaluation for hereditary breast and ovarian cancer syndrome according to the NCCN Guidelines for Genetic/Familial High-Risk Assessment: Breast and Ovarian (3). Referral for genetic assessment for FAMMM syndrome should be considered in a PC patient with a personal or family history of melanoma or multiple dysplastic nevi (261). Patients who meet the aforementioned criteria for LS, PJS, or FAP may also be at increased risk for PC. Consideration for genetic counseling for testing for hereditary pancreatitis is based on expert opinion and warranted for PC patients with a personal history of at least 2 attacks of acute pancreatitis of unknown etiology, a family history of pancreatitis, or early-age onset chronic pancreatitis (262,263).

The risk for development of PC in those patients with an inherited predisposition for PC development is shown in **Table 12**. The risk is highest in hereditary pancreatitis (53-fold risk) (264), PJS (132-fold risk) (197), and FAMMM families with a known *CDKN2A* mutation (13- to 39-fold risk) (265–267). The risk is not as great for *BRCA1* (~2-fold) (268), *BRCA2* (3- to 9-fold) (269,270), LS (9- to 11-fold) (71,271), and *ATM* (~3-fold) (272) mutation carriers based on registry data. There are currently no data available on the risk of developing PC in *PALB2* mutation carriers.

Familial pancreatic cancer (FPC) has been defined by consensus opinion as families with 2 FDRs relatives who do not meet criteria for a known PC-associated hereditary syndrome (262). Case reports first demonstrated families with an excess number of PC cases. Best estimates from case–control and cohort studies suggest that up to 10% of PC patients will have a first- or second-degree relative with PC (273,274). The risk for developing PC in FPC kindreds is dependent on the number of FDRs as shown in **Table 12**. Segregation analysis of families with multiple PC cases has shown an autosomal-dominant inheritance pattern and a 32% lifetime risk for PC development at age 85 years (275). The prospective risk for PC development in individuals with FPC is related to the number of FDRs with PC in the kindred. Those with one or two PC-affected FDRs had a risk ranging from 4- to 7-fold, although whereas those with 3 PC-affected FDRs had a risk ranging from 17 to 32-fold (262,276,277). The complexity in cancer risk assessment has led to the development of a risk prediction model (PancPRO) to provide more detailed risk estimates for individuals from FPC kindreds that take into account the ages at diagnosis, family size, and the relationship between family members (278).

Genetic etiology

Summary statement

- Genetic testing of patients with suspected FPC should include analysis of *BRCA1/2*, *CDKN2A*, *PALB2*, and *ATM*. Evaluation for PJS, LS, and hereditary pancreatitis-associated genes should be considered if other component personal and/or family history criteria are met for the syndrome.

Summary of evidence

In attempts to discover the cause of their PC susceptibility, studies have been performed on familial FPC kindreds for known candidate genes that fall into two groups: (i) genes that cause inherited disorders that are associated with increased risk of PC development (e.g., *BRCA1*, *BRCA2*, and *CDKN2A*) even in the absence of meeting criteria for these hereditary syndromes (279–281) and (ii) recently described genes such as *palladin* (*PALLD*) (282), *ATM* (283), and *PALB2* (284,285) that were discovered by whole genome sequencing or linkage analysis of FPC kindred(s). As shown in **Table 13**, results vary depending on the study population with mutations, for example, in *BRCA1* ranging from 0 to 6% (281,286), *BRCA2* ranging from 0 to 6% (281,287,288), *CDKN2A* ranging from 0 to 20% (280,281,288), and *PALB2* ranging from 0 to 5% (281,285,288). *PALLD* was found to be the susceptibility mutation in a large well-characterized family at the University of Washington (282); however, subsequent studies have not found this gene to be responsible for other FPC families (289,290). Results from a recent study found *ATM* mutations in 2.4% of FPC families (283). Known genetic mutations are responsible for ~20% of the familial clustering of pancreatic cancer. Thus, in the majority of cases, the responsible inherited factor(s) accounting for the increased number of PC cases in these kindreds have not been identified.

Surveillance and management

Recommendations

22. Surveillance of individuals with a genetic predisposition for pancreatic adenocarcinoma should ideally be performed in experienced centers utilizing a multidisciplinary approach and under research conditions. These individuals should be known mutation carriers from hereditary syndromes associated with increased risk of PC (Peutz–Jeghers, hereditary pancreatitis, FAMMM) or members of FPC kindreds with a PC-affected FDR. Because of a lower relative risk for pancreatic adenocarcinoma development in *BRCA1*, *BRCA2*, *PALB2*, *ATM*, and LS families, surveillance should be limited to mutation carriers with a first- or second-degree relative affected with PC (conditional recommendation; very low quality of evidence).
23. Surveillance for PC should be with endoscopic ultrasound and/or MRI of the pancreas annually starting at age 50 years, or 10 years younger than the earliest age of PC in the family. Patients with PJS should start surveillance at age 35 years (conditional recommendation, very low quality of evidence).

Summary of evidence

It is not feasible to screen for PC because of its low incidence in the general population with an estimated lifetime risk in the United States of 1.4% in 2013 (291). Expert opinion has recommended that individuals with a relative risk of more than fivefold when compared with the general population warrant consideration for PC surveillance (73,262,292), with the aim to detect early pancreatic lesions that can be intervened upon. Two such precursor lesions of PC include intraductal papillary mucinous neoplasms and pancreatic intraepithelial neoplasia (293).

Based on this degree of risk, candidates for PC surveillance are limited at this time to unaffected individuals from pancreatic cancer-prone families who are candidates for pancreatic surgery. Unlike colon cancer, in which colonoscopy has been proven to be an effective screening tool to reduce colon cancer-related mortality, there is no proven strategy for PC (294). As stated in a recent International Cancer of the Pancreas Screening Consortium summit (73), recommendations for screening patients with a family history of PC “are primarily based on evidence of increased risk, rather than a proven efficacy of screening.” In light of these acknowledged limitations in PC screening, expert opinion has repeatedly emphasized the importance of performing PC surveillance in the setting of active peer-reviewed research protocols by experienced centers utilizing a multidisciplinary team approach (73,262,292).

Attempts at a unified recommendation for age to commence screening have been unsuccessful with no clear consensus achieved despite being addressed at two international meetings (73,262). Two recently reported surveillance studies have shown that the majority of significant lesions have been found in older patients (295,296). Canto *et al.* (295) reported in their large multicenter surveillance study of high-risk individuals that the prevalence of pancreatic lesions was age related with a significant difference found in the number of lesions detected between patients <50 years old when compared with patients ≥50 years old. Furthermore, all pancreatic lesions with high-grade dysplasia were in patients >65 years of age. Ludwig *et al.* (296) also reported differences in finding a significant abnormality based on age, with a yield of 35% in those >65 years old as compared with only 3% in those ≤65 years old. Several studies in both the hereditary and sporadic setting have found that smokers have an earlier age of PC diagnosis as compared with nonsmokers (297–299), but no data exist regarding whether smoking status should affect surveillance strategy.

Endoscopic ultrasound and MRI/magnetic resonance cholangiopancreatography are the preferred imaging modalities for surveillance as they do not involve the use of radiation like CT scanning and are significantly more accurate in finding pancreatic lesions, particularly small cystic lesions, based on data from CAPS3 study (295).

Recommendation

24. Because of the increased risk for PC development when compared with a pancreatic cyst in the sporadic setting, cystic lesion(s) of the pancreas detected during surveillance of a hereditary pancreatic cancer-prone family member requires evaluation by centers experienced in the care of these high-risk individuals. Determining when surgery is required for pancreatic lesions is difficult and is best individualized after multidisciplinary assessment (conditional recommendation, low quality of evidence).

Summary of evidence

The most common findings in surveillance studies are cystic lesions in the pancreas (295,300). Management of these cysts is unclear as similar to cysts in a nonhereditary setting, most are benign or just have low-grade dysplasia (73,292). Recent consensus recommendations propose that these patients be followed according to international

consensus guidelines for sporadic branch duct intraductal papillary mucinous neoplasms (301), although the majority agreed that surgery should be considered for those branch duct intraductal papillary mucinous neoplasms ≥ 2 cm in size (73).

Prevention strategies—As smoking has been shown to be an independent risk factor for PC in families with FPC, all high-risk individuals should be counseled against smoking (302). In individuals with hereditary pancreatitis, in addition to smoking cessation, a low-fat diet should also be recommended.

Hereditary gastric cancer

Hereditary diffuse gastric cancer

Clinical definitions: *Summary statement*

- Individuals with (i) ≥ 2 cases of diffuse gastric cancer, with at least one diagnosed at <50 years, (ii) ≥ 3 cases of documented diffuse cancer in first- or second-degree relatives independent of age of onset; (iii) diffuse gastric cancer diagnosed at <40 years; and (iv) a personal or family history of diffuse gastric cancer and lobular breast cancer with one diagnosed at <50 years should be evaluated for hereditary diffuse gastric cancer.

Summary of evidence

It is estimated that there were $\sim 21,600$ new cases of gastric cancer in the United States in 2013 (303). The lifetime risk for gastric cancer is $\sim 0.9\%$ (303). The majority of gastric cancer cases are sporadic, but familial clustering is present in $\sim 10\%$ of cases. There are two main types of gastric cancer: diffuse and intestinal. Intestinal-type gastric cancer is a component tumor in LS, familial adenomatous polyposis, and PJS. These syndromes have already been described previously in this guideline, including the risks for gastric cancer, surveillance guidelines for gastric cancer, and genetic testing recommendations, and therefore intestinal forms of gastric cancer will not be discussed in this section.

The only hereditary cancer susceptibility syndrome known to cause diffuse gastric cancer is hereditary diffuse gastric cancer (HDGC). This condition was originally described in three Maori families with autosomal-dominant diffuse gastric cancer in New Zealand in 1964 (304). Approximately 1–3% of diffuse gastric cancers are attributable to HDGC. A genetic evaluation for HDGC is indicated for families having individuals with (i) ≥ 2 cases of diffuse gastric cancer, with at least one diagnosed at <50 years, (ii) ≥ 3 cases of documented diffuse cancer in first- or second-degree relatives independent of age of onset; (iii) diffuse gastric cancer diagnosed at <40 years; and (iv) a personal or family history of diffuse gastric cancer and lobular breast cancer with one diagnosed at <50 years (305).

Genetic etiology

Summary statement

- Genetic testing of individuals who fulfill HDGC clinical criteria should include analysis of *CDH1* mutations.

Summary of the evidence

Mutations in the E-cadherin gene (*CDH1*) were found to be the cause of HDGC in 1998 through linkage analysis (306). The *CDH1* gene encodes for the cell-to-cell adhesion protein E-cadherin. This initial finding led to subsequent studies confirming the presence of *CDH1* mutations in other gastric cancer families (307–309). Genetic testing for the *CDH1* gene is indicated for individuals from families who meet HDGC criteria and should be initiated in an affected individual if at all possible. *CDH1* testing should include sequencing analysis (3–50% of mutations are sequence changes) and deletion/duplication analysis (4% of mutations are large rearrangements) (305). Between 10.5 and 47% of individuals meeting these criteria will be found to have a *CDH1* mutation (305). Once a mutation has been identified in a family, all at-risk individuals should be tested beginning at age 16 years, given the early ages of gastric cancer diagnosis in some families (310).

Surveillance and management

Recommendation

25. Management for patients with HDGC should include (i) prophylactic gastrectomy after age 20 years (>80% risk by age 80 years); (ii) breast cancer surveillance in women beginning at age 35 years with annual mammography and breast MRI and clinical breast examination every 6 months, and (iii) colonoscopy beginning at age 40 years for families that include colon cancer (conditional recommendation, low quality of evidence).

Summary of the evidence

An international consortium study of 11 families with at least 3 cases of diffuse gastric cancer who had tested positive for a *CDH1* gene mutation found that the lifetime risk of developing gastric cancer was 67% for males and 83% for females, with a mean age at diagnosis of 38–40 years of age (range, 14–85) (311). Females also had an increased risk of lobular breast cancer with a lifetime risk of 39% (311). A recent analysis by an international consortium suggests that the risk of gastric cancer is 80% for both men and women, and that the risk of lobular breast cancer is 60% in women (312). There is some evidence that individuals with HDGC are also at increased risk for signet ring cell colon cancer, although exact risk estimates are not known (313). There is a high rate of gastric cancer detection at the time of prophylactic gastrectomy. A recent systematic review (305) found that of 220 previously reported patients who tested positive for *CDH1* mutations, 76.8% underwent a prophylactic gastrectomy whereas 23.2% declined to undergo prophylactic surgery. Among the 169 patients who underwent surgery, 62.7% had negative preoperative endoscopic biopsies and 12.4% (21) tested positive for cancer in their preoperative screening. No information was available on 42 patients. Following gastrectomy, 87% (147) patients had positive histopathology results (including early lesions such as foci of signet ring cells to advanced lesions such as linitis plastica). Only 10% (17 patients) had negative final pathology and it was not reported in 5 patients. It is important to identify both the esophageal and duodenal mucosa at the ends of the surgical specimen because there has been a report of gastric cancer after prophylactic gastrectomy (314).

Gastric cancer surveillance may be used before prophylactic gastrectomy and for patients who decline gastrectomy, but the efficacy is uncertain and should be used with caution. Given the penetrance of this condition, it is recommended that these individuals undergo screening every 6–12 months beginning 5–10 years before the earliest cancer diagnosis in the family (313,314). It is recommended that this is a detailed 30-min endoscopic examination of the gastric mucosa with multiple random biopsies (313,314). Some studies have shown an improved detection rate of early gastric cancer with indigo-carmin staining or a pH-sensitive congo red dye followed by pentagastrin stimulation (315)

Because of limited data, the breast cancer surveillance guidelines are based on those for women with a *BRCA1* or *BRCA2* mutation. Given that lobular breast cancers can be difficult to diagnose by clinical examination and mammography, annual MRI is recommended as part of the surveillance regimen. The colonoscopy guidelines are also based on limited evidence at present.

Informed consent

Genetic testing should only be done in the setting of pre- and posttest genetic counseling. Full informed consent should always be part of the process of cancer genetic counseling. Nongenetics professionals should consider offering cancer genetic testing only if they are able to provide or make available adequate genetic education and counseling as well as access to preventive and surveillance options. Otherwise they should consider referring the patient and family for these services. Standards for informed consent that should be adopted in gastroenterology practices are outlined in **Table 14**.

Emergence of new genetic testing technologies

The advent and commercial availability of next-generation sequencing panels have increased both the complexity and the ease of cancer genetic testing significantly in the past few years. Next-generation sequencing allows for analysis of multiple genes at one time for a lower cost than traditional Sanger sequencing. There are multiple cancer gene panels including anywhere from 6 to 52 genes now offered at a variety of diagnostic laboratories. Some limitations of the new technology include the longer turnaround time (results can take up to 3 months instead of 2–3 weeks) and a higher chance of finding uncertain results (known as variants of uncertain significance) given the large number of genes included. In addition, there are some genes on these panels for which there are very little data available about the associated cancer risks or appropriate management and others for which the cancer risks are so low that the family would be managed based on the cancer history and not the mutation result. In these cases, the results are not useful clinically and can cause confusion for the patient. However, gene panels also may streamline testing for individuals at increased risk for a cancer susceptibility syndrome and reduce greatly the need for sequential tests where multiple diagnoses are under consideration. In general, genetics professionals are using the next-generation sequencing panels for patients with a long list of differential diagnoses (e.g., if testing for >1 possible genetic syndrome) because the panels are much more cost effective and can shorten the diagnostic journey. It is conceivable that gene panels may in a short period of time replace in large part individual genetic tests and

that testing for all the syndromes discussed in this guideline will soon be done simultaneously for at-risk patients.

CONCLUSION

Several well-established hereditary cancer syndromes now exist, each with implications for specific cancer risks in GI and other organ systems. The assessment for cancer susceptibility should be a standard part of patient evaluations in gastroenterology office and endoscopy practices. A systematic and focused family history of cancer and premalignant conditions is the first step and sufficient to screen for ~ 10–15% of patients who may need more detailed risk assessment by more extensive family history assessment, genetic counseling, and genetic testing. The underlying genetic etiologies for several syndromes are now well established, and the array of cancer susceptibility genes is continually expanding. Genetic testing is widely available and should be part of standard of care of patients at increased risk for a hereditary cancer syndrome. Mutation carriers and at-risk individuals require intensive surveillance, possibly prophylactic surgery, and family counseling, and management needs to be individualized based on the syndrome under consideration, as well as the specifics of the family history at hand. There is a dire need of organized collaborative international efforts to study benefits of surveillance and surgical strategies in patients with these relatively rare syndromes in order to be able to offer truly evidence-based management recommendations.

ACKNOWLEDGMENTS

This guideline was produced in collaboration with the Practice Parameters Committee of the American College of Gastroenterology. The Committee gives special thanks to Christine Y. Hachem, MD, FACG, who served as guideline monitor for this document. We are grateful to Christine Hachem, Lauren Gerson, and Maria Susano for assistance with guideline development, to Paul Moayyedi for reviewing and grading recommendations, and to Chinedu Ukaegbu for help with drafting of the manuscript tables and assembling reference libraries.

Financial support: This study was supported by the National Cancer Institute K24-11311 (to S.S.).

REFERENCES

1. Guyatt GH, Oxman AD, Vist GE, et al. GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. *BMJ*. 2008; 336:924. [PubMed: 18436948]
2. Lu KH, Wood ME, Daniels M, et al. American Society of Clinical Oncology Expert Statement: collection and use of a cancer family history for oncology providers. *J Clin Oncol*. 2014; 32:833–40. [PubMed: 24493721]
3. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Genetic/Familial High-Risk Assessment: Breast and Ovarian. Version 4.2013. © National Comprehensive Cancer Network, Inc 2014. All rights reserved. [cited 2014 March 13]; To view the most recent and complete version of the guidelines, go online to [NCCN.org](http://www.nccn.org). NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, NCCN GUIDELINES®, and all other NCCN Content are trademarks owned by the National Comprehensive Cancer Network, Inc]. Available from http://www.nccn.org/professionals/physician_gls/f_guidelines.asp.
4. Ziogas A, Anton-Culver H. Validation of family history data in cancer family registries. *Am J Prev Med*. 2003; 24:190–8. [PubMed: 12568826]
5. Wideroff L, Garceau AO, Greene MH, et al. Coherence and completeness of population-based family cancer reports. *Cancer Epidemiol Biomarkers Prev*. 2010; 19:799–810. [PubMed: 20160272]

6. Aaltonen LA, Salovaara R, Kristo P, et al. Incidence of hereditary nonpolyposis colorectal cancer and the feasibility of molecular screening for the disease. *N Engl J Med.* 1998; 338:1481–7. [PubMed: 9593786]
7. Barnetson RA, Tenesa A, Farrington SM, et al. Identification and survival of carriers of mutations in DNA mismatch-repair genes in colon cancer. *N Engl J Med.* 2006; 354:2751–63. [PubMed: 16807412]
8. Hampel H, Frankel WL, Martin E, et al. Screening for the Lynch syndrome (hereditary nonpolyposis colorectal cancer). *N Engl J Med.* 2005; 352:1851–60. [PubMed: 15872200]
9. Piñol V, Castells A, Andreu M, et al. Accuracy of revised Bethesda guidelines, microsatellite instability, and immunohistochemistry for the identification of patients with hereditary nonpolyposis colorectal cancer. *JAMA.* 2005; 293:1986–94. [PubMed: 15855432]
10. Salovaara R, Loukola A, Kristo P, et al. Population-based molecular detection of hereditary nonpolyposis colorectal cancer. *J Clin Oncol.* 2000; 18:2193–200. [PubMed: 10829038]
11. Berg A, Armstrong K, Botkin J, et al. Recommendations from the EGAPP Working Group: genetic testing strategies in newly diagnosed individuals with colorectal cancer aimed at reducing morbidity and mortality from Lynch syndrome in relatives. *Genet Med.* 2009; 11:35–41. [PubMed: 19125126]
12. Hampel H, Frankel WL, Martin E, et al. Feasibility of screening for Lynch syndrome among patients with colorectal cancer. *J Clin Oncol.* 2008; 26:5783–8. [PubMed: 18809606]
13. Akiyama Y, Sato H, Yamada T, et al. Germ-line mutation of the hMSH6/ GTBP gene in an atypical hereditary nonpolyposis colorectal cancer kindred. *Cancer Res.* 1997; 57:3920–3. [PubMed: 9307272]
14. Bronner CE, Baker SM, Morrison PT, et al. Mutation in the DNA mismatch repair gene homologue hMLH1 is associated with hereditary non-polyposis colon cancer. *Nature.* 1994; 368:258–61. [PubMed: 8145827]
15. Fishel R, Lescoe MK, Rao MR, et al. The human mutator gene homolog MSH2 and its association with hereditary nonpolyposis colon cancer. *Cell.* 1993; 75:1027–38. [PubMed: 8252616]
16. Leach FS, Nicolaidis NC, Papadopoulos N, et al. Mutations of a mutS homolog in hereditary nonpolyposis colorectal cancer. *Cell.* 1993; 75:1215–25. [PubMed: 8261515]
17. Ligtenberg MJ, Kuiper RP, Chan TL, et al. Heritable somatic methylation and inactivation of MSH2 in families with Lynch syndrome due to deletion of the 3' exons of TACSTD1. *Nat Genet.* 2009; 41:112–7. [PubMed: 19098912]
18. Lindblom A, Tannergard P, Werelius B, et al. Genetic mapping of a second locus predisposing to hereditary non-polyposis colon cancer. *Nat Genet.* 1993; 5:279–82. [PubMed: 7903889]
19. Liu B, Parsons RE, Hamilton SR, et al. hMSH2 mutations in hereditary nonpolyposis colorectal cancer kindreds. *Cancer Res.* 1994; 54:4590–4. [PubMed: 8062247]
20. Miyaki M, Konishi M, Tanaka K, et al. Germline mutation of MSH6 as the cause of hereditary nonpolyposis colorectal cancer. *Nat Genet.* 1997; 17:271–2. 1997. [PubMed: 9354786]
21. Nicolaidis NC, Papadopoulos N, Liu B, et al. Mutations of two PMS homologues in hereditary nonpolyposis colon cancer. *Nature.* 1994; 371:75–80. [PubMed: 8072530]
22. Papadopoulos N, Nicolaidis NC, Wei YF, et al. Mutation of a mutL homolog in hereditary colon cancer. *Science.* 1994; 263:1625–9. [PubMed: 8128251]
23. Peltomaki P, Aaltonen LA, Sistonen P, et al. Genetic mapping of a locus predisposing to human colorectal cancer. *Science.* 1993; 260:810–2. [PubMed: 8484120]
24. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Genetic/Familial High-Risk Assessment: Colorectal V.1.2014. © National Comprehensive Cancer Network, Inc 2014. All rights reserved. [cited 2014 March 13]; To view the most recent and complete version of the guideline, go online to NCCN.org. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, NCCN GUIDELINES®, and all other NCCN Content are trademarks owned by the National Comprehensive Cancer Network, Inc]. Available from http://www.nccn.org/professionals/physician_gls/f_guidelines.asp.
25. Giardiello FM, Allen JI, Axilbund JE, et al. Guidelines on genetic evaluation and management of Lynch syndrome: a consensus statement by the US Multi-society Task Force on colorectal cancer. *Am J Gastroenterol.* 2014; 109:1159–79. [PubMed: 25070057]

26. Dinh TA, Rosner BI, Atwood JC, et al. Health benefits and cost-effectiveness of primary genetic screening for Lynch syndrome in the general population. *Cancer Prev Res.* 2011; 4:9–22.
27. Kastrinos F, Allen JI, Stockwell DH, et al. Development and validation of a colon cancer risk assessment tool for patients undergoing colonoscopy. *Am J Gastroenterol.* 2009; 104:1508–18. [PubMed: 19491864]
28. Jass J, Stewart S, Stewart J, et al. Hereditary non-polyposis colorectal cancer--morphologies, genes and mutations. *Mutat Res.* 1994; 310:125–33. [PubMed: 7523876]
29. Edelstein DL, Axilbund J, Baxter M, et al. Rapid development of colorectal neoplasia in patients with Lynch syndrome. *Clin Gastroenterol Hepatol.* 2011; 9:340–3. [PubMed: 21070872]
30. Alarcon F, Lasset C, Carayol J, et al. Estimating cancer risk in HNPCC by the GRL method. *Eur J Hum Genet.* 2007; 15:831–6. [PubMed: 17473834]
31. Bonadona V, Bonaïti B, Olschwang S, et al. Cancer risks associated with germline mutations in MLH1, MSH2, and MSH6 genes in Lynch syndrome. *JAMA.* 2011; 305:2304. [PubMed: 21642682]
32. Choi YH, Cotterchio M, McKeown-Eyssen G, et al. Penetrance of colorectal cancer among MLH1/MSH2 carriers participating in the colorectal cancer familial registry in Ontario. *Heredit Cancer Clin Pract.* 2009; 7:1897–4287.
33. Dunlop MG, Farrington SM, Carothers AD, et al. Cancer risk associated with germline DNA mismatch repair gene mutations. *Hum Mol Genet.* 1997; 6:105–10. [PubMed: 9002677]
34. Jenkins MA, Baglietto L, Dowty JG, et al. Cancer risks for mismatch repair gene mutation carriers: a population-based early onset case-family study. *Clin Gastroenterol Hepatol.* 2006; 4:489–98. [PubMed: 16616355]
35. Quehenberger F, Vasen HF, van Houwelingen HC. Risk of colorectal and endometrial cancer for carriers of mutations of the hMLH1 and hMSH2 gene: correction for ascertainment. *J Med Genet.* 2005; 42:491–6. [PubMed: 15937084]
36. Hendriks Y, Wagner A, Morreau H, et al. Cancer risk in hereditary nonpolyposis colorectal cancer due to MSH6 mutations: impact on counseling and surveillance. *Gastroenterology.* 2004; 127:17–25. [PubMed: 15236168]
37. Senter L, Clendenning M, Sotamaa K, et al. The clinical phenotype of Lynch syndrome due to germ-line PMS2 mutations. *Gastroenterology.* 2008; 135:419–28. [PubMed: 18602922]
38. Hampel H, Stephens JA, Pukkala E, et al. Cancer risk in hereditary nonpolyposis colorectal cancer syndrome: later age of onset. *Gastroenterology.* 2005; 129:415–21. [PubMed: 16083698]
39. Vasen H. Clinical description of the Lynch syndrome [hereditary nonpolyposis colorectal cancer (HNPCC)]. *Fam Cancer.* 2005; 4:219–25. [PubMed: 16136381]
40. Howlader, N.; Noone, A.; Krapcho, M., et al. SEER Cancer Statistics Review, 1975–2009 (Vintage 2009 Populations). National Cancer Institute; Bethesda, MA: 2012.
41. Järvinen HJ, Aarnio M, Mustonen H, et al. Controlled 15-year trial on screening for colorectal cancer in families with hereditary nonpolyposis colorectal cancer. *Gastroenterology.* 2000; 118:829–34. [PubMed: 10784581]
42. Järvinen HJ, Mecklin J-P, Sistonen P. Screening reduces colorectal cancer rate in families with hereditary nonpolyposis colorectal cancer. *Gastroenterology.* 1995; 108:1405–11. [PubMed: 7729632]
43. Järvinen HJ, Renkonen-Sinisalo L, Aktán-Collán K, et al. Ten years after mutation testing for Lynch syndrome: cancer incidence and outcome in mutation-positive and mutation-negative family members. *J Clin Oncol.* 2009; 27:4793–7. [PubMed: 19720893]
44. Dove-Edwin I, Sasieni P, Adams J, et al. Prevention of colorectal cancer by colonoscopic surveillance in individuals with a family history of colorectal cancer: 16 year, prospective, follow-up study. *BMJ.* 2005; 331:1047. [PubMed: 16243849]
45. Engel C, Rahner N, Schulmann K, et al. Efficacy of annual colonoscopic surveillance in individuals with hereditary nonpolyposis colorectal cancer. *Clin Gastroenterol Hepatol.* 2010; 8:174–82. [PubMed: 19835992]
46. Stuckless S, Green J, Morgenstern M, et al. Impact of colonoscopic screening in male and female Lynch syndrome carriers with an MSH2 mutation. *Clin Genet.* 2012; 82:439–45. [PubMed: 22011075]

47. Vasen HF, Abdirahman M, Brohet R, et al. One to 2-year surveillance intervals reduce risk of colorectal cancer in families with Lynch syndrome. *Gastroenterology*. 2010; 138:2300–6. [PubMed: 20206180]
48. de Vos tot Nederveen Cappel W, Nagengast FM, Griffioen G, et al. Surveillance for hereditary nonpolyposis colorectal cancer: a long-term study of 114 families. *Dis Colon Rectum*. 2002; 45:1588–94. [PubMed: 12473880]
49. Baglietto L, Lindor NM, Dowty JG, et al. Risks of Lynch syndrome cancers for MSH6 mutation carriers. *J Natl Cancer Inst*. 2010; 102:193–201. [PubMed: 20028993]
50. Parry S, Win AK, Parry B, et al. Metachronous colorectal cancer risk for mismatch repair gene mutation carriers: the advantage of more extensive colon surgery. *Gut*. 2011; 60:950–7. [PubMed: 21193451]
51. Win A, Parry S, Parry B, et al. Risk of metachronous colon cancer following surgery for rectal cancer in mismatch repair gene mutation carriers. *Ann Surg Oncol*. 2013; 20:1829–36. [PubMed: 23358792]
52. Haanstra JF, de Vos Tot Nederveen Cappel WH, Gopie JP, et al. Quality of life after surgery for colon cancer in patients with Lynch syndrome: partial versus subtotal colectomy. *Dis Colon Rectum*. 2012; 55:653–9. [PubMed: 22595844]
53. Barrow E, Robinson L, Alduaij W, et al. Cumulative lifetime incidence of extracolonic cancers in Lynch syndrome: a report of 121 families with proven mutations. *Clin Genet*. 2009; 75:141–9. [PubMed: 19215248]
54. Engel C, Loeffler M, Steinke V, et al. Risks of less common cancers in proven mutation carriers with Lynch syndrome. *J Clin Oncol*. 2012; 30:4409–15. [PubMed: 23091106]
55. Watson P, Vasen HF, Mecklin JP, et al. The risk of extra-colonic, extra-endometrial cancer in the Lynch syndrome. *Int J Cancer*. 2008; 123:444–9. [PubMed: 18398828]
56. Dove-Edwin I, Boks D, Goff S, et al. The outcome of endometrial carcinoma surveillance by ultrasound scan in women at risk of hereditary nonpolyposis colorectal carcinoma and familial colorectal carcinoma. *Cancer*. 2002; 94:1708–12. [PubMed: 11920532]
57. Renkonen-Sinisalo L, Bützow R, Leminen A, et al. Surveillance for endometrial cancer in hereditary nonpolyposis colorectal cancer syndrome. *Int J Cancer*. 2007; 120:821–4. [PubMed: 17096354]
58. Schmeler KM, Lynch HT, Chen L-m, et al. Prophylactic surgery to reduce the risk of gynecologic cancers in the Lynch syndrome. *N Engl J Med*. 2006; 354:261–9. [PubMed: 16421367]
59. Yang KY, Caughey AB, Little SE, et al. A cost-effectiveness analysis of prophylactic surgery versus gynecologic surveillance for women from hereditary non-polyposis colorectal cancer (HNPCC) Families. *Fam Cancer*. 2011; 10:535–43. [PubMed: 21538078]
60. Kwon J, Sun C, Peterson S, et al. Cost-effectiveness analysis of prevention strategies for gynecologic cancers in Lynch syndrome. *Cancer*. 2008; 113:326–35. [PubMed: 18506736]
61. Capelle LG, Van Grieken NC, Lingsma HF, et al. Risk and epidemiological time trends of gastric cancer in Lynch syndrome carriers in the Netherlands. *Gastroenterology*. 2010; 138:487–92. [PubMed: 19900449]
62. Aarnio M, Salovaara R, Aaltonen LA, et al. Features of gastric cancer in hereditary non-polyposis colorectal cancer syndrome. *Int J Cancer*. 1997; 74:551–5. [PubMed: 9355980]
63. Aarnio M, Sankila R, Pukkala E, et al. Cancer risk in mutation carriers of DNA-mismatch-repair genes. *Int J Cancer*. 1999; 81:214–8. [PubMed: 10188721]
64. Vasen H, Stormorken A, Menko F, et al. MSH2 mutation carriers are at higher risk of cancer than MLH1 mutation carriers: a study of hereditary nonpolyposis colorectal cancer families. *J Clin Oncol*. 2001; 19:4074–80. [PubMed: 11600610]
65. Schulmann K, Brasch FE, Kunstmann E, et al. HNPCC-associated small bowel cancer: clinical and molecular characteristics. *Gastroenterology*. 2005; 128:590–9. [PubMed: 15765394]
66. Saurin J-C, Pilleul F, Soussan E, et al. Small-bowel capsule endoscopy diagnoses early and advanced neoplasms in asymptomatic patients with Lynch syndrome. *Endoscopy*. 2010; 42:1057–62. [PubMed: 20821360]

67. Van der Post R, Kiemeny L, Ligtenberg M, et al. Risk of urothelial bladder cancer in Lynch syndrome is increased, in particular among MSH2 mutation carriers. *J Med Genet.* 2010; 47:464–70. [PubMed: 20591884]
68. Myrhøj T, Andersen M-B, Bernstein I. Screening for urinary tract cancer with urine cytology in Lynch syndrome and familial colorectal cancer. *Fam Cancer.* 2008; 7:303–7. [PubMed: 18389386]
69. USPSTF. US Preventive Services Task Force. Guide to Clinical Preventive Services-17.. Screening for Bladder Cancer. 1996. [cited; 2nd Edition [181-186] Available from <http://odphp.osophs.dhhs.gov/pubs/guidecps/PDF/CH17.PDF>
70. Thériault GP, Tremblay CG, Armstrong BG. Bladder cancer screening among primary aluminum production workers in Quebec. *J Occup Environ Med.* 1990; 32:869.
71. Win AK, Young JP, Lindor NM, et al. Colorectal and other cancer risks for carriers and noncarriers from families with a DNA mismatch repair gene mutation: a prospective cohort study. *J Clin Oncol.* 2012; 30:958–64. [PubMed: 22331944]
72. Kastros F, Stoffel EM, Balmaña J, et al. Phenotype comparison of MLH1 and MSH2 mutation carriers in a cohort of 1,914 individuals undergoing clinical genetic testing in the United States. *Cancer Epidemiol Biomark Prev.* 2008; 17:2044–51.
73. Canto MI, Harinck F, Hruban RH, et al. International Cancer of the Pancreas Screening (CAPS) Consortium summit on the management of patients with increased risk for familial pancreatic cancer. *Gut.* 2013; 62:339–47. [PubMed: 23135763]
74. Müller A, Edmonston TB, Corao DA, et al. Exclusion of breast cancer as an integral tumor of hereditary nonpolyposis colorectal cancer. *Cancer Res.* 2002; 62:1014–9. [PubMed: 11861375]
75. Vasen H, Morreau H, Nortier J. Is breast cancer part of the tumor spectrum of hereditary nonpolyposis colorectal cancer? *Am J Hum Genet.* 2001; 68:1533. [PubMed: 11359214]
76. Raymond VM, Mukherjee B, Wang F, et al. Elevated risk of prostate cancer among men with Lynch syndrome. *J Clin Oncol.* 2013; 31:1713–8. [PubMed: 23530095]
77. Winkels RM, Botma A, Van Duijnhoven FJ, et al. Smoking increases the risk for colorectal adenomas in patients with Lynch syndrome. *Gastroenterology.* 2012; 142:241–7. [PubMed: 22062356]
78. Botma A, Nagengast FM, Braem MG, et al. Body mass index increases risk of colorectal adenomas in men with Lynch syndrome: the GEOLynch cohort study. *J Clin Oncol.* 2010; 28:4346–53. [PubMed: 20733131]
79. Mathers JC, Movahedi M, Macrae F, et al. Long-term effect of resistant starch on cancer risk in carriers of hereditary colorectal cancer: an analysis from the CAPP2 randomised controlled trial. *Lancet Oncol.* 2012; 13:1242–9. [PubMed: 23140761]
80. Burn J, Gerdes A-M, Macrae F, et al. Long-term effect of aspirin on cancer risk in carriers of hereditary colorectal cancer: an analysis from the CAPP2 randomised controlled trial. *Lancet.* 2012; 378:2081–7. [PubMed: 22036019]
81. Bussey, HJR. *Family Studies, Histopathology, Differential Diagnosis and Results of Treatment.* Johns Hopkins University Press; Baltimore: 1975. *Familial Polyposis Coli.*
82. Jarvinen HJ. Epidemiology of familial adenomatous polyposis in Finland: impact of family screening on the colorectal cancer rate and survival. *Gut.* 1992; 33:357–60. [PubMed: 1314763]
83. Bulow S, Faurschou Nielsen T, Bulow C, et al. The incidence rate of familial adenomatous polyposis. Results from the Danish Polyposis Register. *Int J Colorectal Dis.* 1996; 11:88–91. [PubMed: 8739833]
84. Bisgaard ML, Fenger K, Bulow S, et al. Familial adenomatous polyposis (FAP): frequency, penetrance, and mutation rate. *Hum Mutat.* 1994; 3:121–5. [PubMed: 8199592]
85. Bjork J, Akerbrant H, Iselius L, et al. Epidemiology of familial adenomatous polyposis in Sweden: changes over time and differences in phenotype between males and females. *Scand J Gastroenterol.* 1999; 34:1230–5. [PubMed: 10636071]
86. Iwama T, Tamura K, Morita T, et al. A clinical overview of familial adenomatous polyposis derived from the database of the Polyposis Registry of Japan. *Int J Clin Oncol.* 2004; 9:308–16. [PubMed: 15375708]

87. Scheuner MT, McNeel TS, Freedman AN. Population prevalence of familial cancer and common hereditary cancer syndromes. The 2005 California Health Interview Survey. *Genet Med*. 2010; 12:726–35. [PubMed: 20921897]
88. Hernegger GS, Moore HG, Guillem JG. Attenuated familial adenomatous polyposis: an evolving and poorly understood entity. *Dis Colon Rectum*. 2002; 45:127–34. discussion 134-6. [PubMed: 11786778]
89. Knudsen AL, Bisgaard ML, Bulow S. Attenuated familial adenomatous polyposis (AFAP). A review of the literature. *Fam Cancer*. 2003; 2:43–55. [PubMed: 14574166]
90. Burt RW, Leppert MF, Slattery ML, et al. Genetic testing and phenotype in a large kindred with attenuated familial adenomatous polyposis. *Gastroenterology*. 2004; 127:444–51. [PubMed: 15300576]
91. Al-Tassan N, Chmiel NH, Maynard J, et al. Inherited variants of MYH associated with somatic G:C-->T:A mutations in colorectal tumors. *Nat Genet*. 2002; 30:227–32. [PubMed: 11818965]
92. Perchiniak EM, Groden J. Mechanisms regulating microtubule binding, DNA replication, and apoptosis are controlled by the intestinal tumor suppressor APC. *Curr Colorectal Cancer Rep*. 2011; 7:145–51. [PubMed: 23308069]
93. Burt R, Neklason DW. Genetic testing for inherited colon cancer. *Gastroenterology*. 2005; 128:1696–716. [PubMed: 15887160]
94. Aretz S, Uhlhaas S, Caspari R, et al. Frequency and parental origin of de novo APC mutations in familial adenomatous polyposis. *Eur J Hum Genet*. 2004; 12:52–8. [PubMed: 14523376]
95. Aretz S, Stienen D, Friedrichs N, et al. Somatic APC mosaicism: a frequent cause of familial adenomatous polyposis (FAP). *Hum Mutat*. 2007; 28:985–92. [PubMed: 17486639]
96. Hes FJ, Nielsen M, Bik EC, et al. Somatic APC mosaicism: an underestimated cause of polyposis coli. *Gut*. 2008; 57:71–6. [PubMed: 17604324]
97. Schwab AL, Tuohy TM, Condie M, et al. Gonadal mosaicism and familial adenomatous polyposis. *Fam Cancer*. 2008; 7:173–7. [PubMed: 18026870]
98. Tuohy TM, Burt RW. Somatic mosaicism: a cause for unexplained cases of FAP? *Gut*. 2008; 57:10–12. [PubMed: 18094199]
99. Chung DC. The genetic basis of colorectal cancer: insights into critical pathways of tumorigenesis. *Gastroenterology*. 2000; 119:854–65. [PubMed: 10982779]
100. Nagase H, Nakamura Y. Mutations of the APC (adenomatous polyposis coli) gene. *Hum Mutat*. 1993; 2:425–34. [PubMed: 8111410]
101. Groves C, Lamlum H, Crabtree M, et al. Mutation cluster region, association between germline and somatic mutations and genotype-phenotype correlation in upper gastrointestinal familial adenomatous polyposis. *Am J Pathol*. 2002; 160:2055–61. [PubMed: 12057910]
102. Nieuwenhuis MH, Vasen HF. Correlations between mutation site in APC and phenotype of familial adenomatous polyposis (FAP): a review of the literature. *Crit Rev Oncol Hematol*. 2007; 61:153–61. [PubMed: 17064931]
103. Nielsen M, Bik E, Hes FJ, et al. Genotype-phenotype correlations in 19 Dutch cases with APC gene deletions and a literature review. *Eur J Hum Genet*. 2007; 15:1034–42. [PubMed: 17568392]
104. Sieber OM, Lipton L, Crabtree M, et al. Multiple colorectal adenomas, classic adenomatous polyposis, and germ-line mutations in MYH. *N Engl J Med*. 2003; 348:791–9. [PubMed: 12606733]
105. Lipton L, Tomlinson I. The multiple colorectal adenoma phenotype and MYH, a base excision repair gene. *Clin Gastroenterol Hepatol*. 2004; 2:633–8. [PubMed: 15290654]
106. Halford SE, Rowan AJ, Lipton L, et al. Germline mutations but not somatic changes at the MYH locus contribute to the pathogenesis of unselected colorectal cancers. *Am J Pathol*. 2003; 162:1545–8. [PubMed: 12707038]
107. Nielsen M, Morreau H, Vasen HF, et al. MUTYH-associated polyposis (MAP). *Crit Rev Oncol Hematol*. 2011; 79:1–16. [PubMed: 20663686]
108. Jones S, Emmerson P, Maynard J, et al. Biallelic germline mutations in MYH predispose to multiple colorectal adenoma and somatic G:C-->T:A mutations. *Hum Mol Genet*. 2002; 11:2961–7. [PubMed: 12393807]

109. Sampson JR, Dolwani S, Jones S, et al. Autosomal recessive colorectal adenomatous polyposis due to inherited mutations of MYH. *The Lancet*. 2003; 362:39–41.
110. Enholm S, Hienonen T, Suomalainen A, et al. Proportion and phenotype of MYH-associated colorectal neoplasia in a population-based series of Finnish colorectal cancer patients. *Am J Pathol*. 2003; 163:827–32. [PubMed: 12937124]
111. Filipe B, Baltazar C, Albuquerque C, et al. APC or MUTYH mutations account for the majority of clinically well-characterized families with FAP and AFAP phenotype and patients with more than 30 adenomas. *Clin Genet*. 2009; 76:242–55. [PubMed: 19793053]
112. Wang L, Baudhuin LM, Boardman LA, et al. MYH mutations in patients with attenuated and classic polyposis and with young-onset colorectal cancer without polyps. *Gastroenterology*. 2004; 127:9–16. [PubMed: 15236166]
113. Jo W-S, Bandipalliam P, Shannon KM, et al. Correlation of polyp number and family history of colon cancer with germline MYH mutations. *Clin Gastroenterol Hepatol*. 2005; 3:1022–8. [PubMed: 16234049]
114. Venesio T, Molatore S, Cattaneo F, et al. High frequency of MYH gene mutations in a subset of patients with familial adenomatous polyposis. *Gastroenterology*. 2004; 126:1681–5. [PubMed: 15188161]
115. Knopperts AP, Nielsen M, Niessen RC, et al. Contribution of bi-allelic germline MUTYH mutations to early-onset and familial colorectal cancer and to low number of adenomatous polyps: case-series and literature review. *Fam Cancer*. 2013; 12:43–50. [PubMed: 23007840]
116. Farrington SM, Tenesa A, Barnetson R, et al. Germline susceptibility to colorectal cancer because of base-excision repair gene defects. *Am J Hum Genet*. 2005; 77:112–9. [PubMed: 15931596]
117. Palles C, Cazier JB, Howarth KM, et al. Germline mutations affecting the proofreading domains of POLE and POLD1 predispose to colorectal adenomas and carcinomas. *Nat Genet*. 2012; 45:136–44. [PubMed: 23263490]
118. Valle L, Hernández-Illán E, Bellido F, et al. New insights into POLE and POLD1 germline mutations in familial colorectal cancer and polyposis. *Hum Mol Genet*. 2014; 23:3506–12. [PubMed: 24501277]
119. Petersen GM, Slack J, Nakamura Y. Screening guidelines and premorbid diagnosis of familial adenomatous polyposis using linkage. *Gastroenterology*. 1991; 100:1658–64. [PubMed: 1673441]
120. Giardiello FM, Krush AJ, Petersen GM, et al. Phenotypic variability of familial adenomatous polyposis in 11 unrelated families with identical APC gene mutation. *Gastroenterology*. 1994; 106:1542–7. [PubMed: 8194700]
121. Roncucci L, Stamp D, Medline A, et al. Identification and quantification of aberrant crypt foci and microadenomas in the human colon. *Hum Pathol*. 1991; 22:287–94. [PubMed: 1706308]
122. Jang YS, Steinhagen RM, Heimann TM. Colorectal cancer in familial adenomatous polyposis. *Dis Colon Rectum*. 1997; 40:312–6. [PubMed: 9118746]
123. Giardiello FM, Brensinger JD, Petersen GM. AGA technical review on hereditary colorectal cancer and genetic testing. *Gastroenterology*. 2001; 121:198–213. [PubMed: 11438509]
124. Burt RW. Colon cancer screening. *Gastroenterology*. 2000; 119:837–53. [PubMed: 10982778]
125. Guillem JG, Smith AJ, Culle J, et al. Gastrointestinal polyposis syndromes. *Curr Probl Surg*. 1999; 36:217–323. [PubMed: 10212621]
126. Jasperson KW, Tuohy TM, Neklason DW, et al. Hereditary and familial colon cancer. *Gastroenterology*. 2010; 138:2044–58. [PubMed: 20420945]
127. Vasen HF, Moslein G, Alonso A, et al. Guidelines for the clinical management of familial adenomatous polyposis (FAP). *Gut*. 2008; 57:704–13. [PubMed: 18194984]
128. Bjork JA, Akerbrant HI, Iselius LE, et al. Risk factors for rectal cancer morbidity and mortality in patients with familial adenomatous polyposis after colectomy and ileorectal anastomosis. *Dis Colon Rectum*. 2000; 43:1719–25. [PubMed: 11156457]
129. Bulow S. Results of national registration of familial adenomatous polyposis. *Gut*. 2003; 52:742–6. [PubMed: 12692062]
130. Heiskanen I, Luostarinen T, Jarvinen HJ. Impact of screening examinations on survival in familial adenomatous polyposis. *Scand J Gastroenterol*. 2000; 35:1284–7. [PubMed: 11199368]

131. Gibbons DC, Sinha A, Phillips RK, et al. Colorectal cancer: no longer the issue in familial adenomatous polyposis? *Fam Cancer*. 2011; 10:11–20. [PubMed: 21052851]
132. Friedl W, Caspari R, Sengteller M, et al. Can APC mutation analysis contribute to therapeutic decisions in familial adenomatous polyposis? Experience from 680 FAP families. *Gut*. 2001; 48:515–21. [PubMed: 11247896]
133. Cleary SP, Cotterchio M, Jenkins MA, et al. Germline MutY human homo-logue mutations and colorectal cancer: a multisite case-control study. *Gastroenterology*. 2009; 136:1251–60. [PubMed: 19245865]
134. Nielsen M, Joerink-van de Beld MC, Jones N, et al. Analysis of MUTYH genotypes and colorectal phenotypes in patients with MUTYH-associated polyposis. *Gastroenterology*. 2009; 136:471–6. [PubMed: 19032956]
135. Lubbe SJ, Di Bernardo MC, Chandler IP, et al. Clinical implications of the colorectal cancer risk associated with MUTYH mutation. *J Clin Oncol*. 2009; 27:3975–80. [PubMed: 19620482]
136. Boparai KS, Dekker E, Van Eeden S, et al. Hyperplastic polyps and sessile serrated adenomas as a phenotypic expression of MYH-associated polyposis. *Gastroenterology*. 2008; 135:2014–8. [PubMed: 19013464]
137. da Luz Moreira A, Church JM, Burke CA. The evolution of prophylactic colorectal surgery for familial adenomatous polyposis. *Dis Colon Rectum*. 2009; 52:1481–6. [PubMed: 19617764]
138. Galiatsatos P, Foulkes WD. Familial adenomatous polyposis. *Am J Gastroenterol*. 2006; 101:385–98. [PubMed: 16454848]
139. Nieuwenhuis MH, Bulow S, Bjork J, et al. Genotype predicting phenotype in familial adenomatous polyposis: a practical application to the choice of surgery. *Dis Colon Rectum*. 2009; 52:1259–63. [PubMed: 19571702]
140. Kartheuser A, Stangherlin P, Brandt D, et al. Restorative proctocolectomy and ileal pouch-anal anastomosis for familial adenomatous polyposis revisited. *Fam Cancer*. 2006; 5:241–60. [PubMed: 16998670]
141. Saurin JC, Napoleon B, Gay G, et al. Endoscopic management of patients with familial adenomatous polyposis (FAP) following a colectomy. *Endoscopy*. 2005; 37:499–501.
142. Smith JC, Schaffer MW, Ballard BR, et al. Adenocarcinomas after prophylactic surgery for familial adenomatous polyposis. *J Cancer Ther*. 2013; 4:260–70. [PubMed: 23875116]
143. Ooi BS, Remzi FH, Gramlich T, et al. Anal transitional zone cancer after restorative proctocolectomy and ileoanal anastomosis in familial adenomatous polyposis: report of two cases. *Dis Colon Rectum*. 2003; 46:1418–23. discussion 1422-3. [PubMed: 14530685]
144. Bulow S, Bjork J, Christensen IJ, et al. Duodenal adenomatosis in familial adenomatous polyposis. *Gut*. 2004; 53:381–6. [PubMed: 14960520]
145. Lopez-Ceron M, van den Broek FJ, Mathus-Vliegen EM, et al. The role of high-resolution endoscopy and narrow-band imaging in the evaluation of upper GI neoplasia in familial adenomatous polyposis. *Gastrointest Endosc*. 2013; 77:542–50. [PubMed: 23352497]
146. Bjork J, Akerbrant H, Iselius L, et al. Periampullary adenomas and adenocarcinomas in familial adenomatous polyposis: cumulative risks and APC gene mutations. *Gastroenterology*. 2001; 121:1127–35. [PubMed: 11677205]
147. Latchford AR, Neale KF, Spigelman AD, et al. Features of duodenal cancer in patients with familial adenomatous polyposis. *Clin Gastroenterol Hepatol*. 2009; 7:659–63. [PubMed: 19281862]
148. Drini M, Speer A, Dow C, et al. Management of duodenal adenomatosis in FAP: single centre experience. *Fam Cancer*. 2012; 11:167–73. [PubMed: 22131053]
149. Iaquinto G, Fornasari M, Quaia M, et al. Capsule endoscopy is useful and safe for small-bowel surveillance in familial adenomatous polyposis. *Gastrointest Endosc*. 2008; 67:61–7. [PubMed: 18155426]
150. Tulchinsky H, Keidar A, Strul H, et al. Extracolonic manifestations of familial adenomatous polyposis after proctocolectomy. *Arch Surg*. 2005; 140:159–63. discussion 164. [PubMed: 15723997]

151. Spigelman, AD.; Phillips, RKS. The upper gastrointestinal tract.. In: Phillips, RKS.; Spigelman, AD.; Thomason, JPS., editors. *Familial Adenomatous Polyposis and Other Syndromes* Edward Arnold. London: 1994. p. 106
152. Tytgat GN. Surveillance of familial adenomatous polyposis patients after ileorectal anastomosis or ileoanal pouch anastomosis. *Gastrointest Endosc Clin N Am.* 1997; 7:111–27. [PubMed: 8995117]
153. Alderlieste YA, Rauws EA, Mathus-Vliegen EM, et al. Prospective enteroscopic evaluation of jejunal polyposis in patients with familial adenomatous polyposis and advanced duodenal polyposis. *Fam Cancer.* 2013; 12:51–6. [PubMed: 23054214]
154. Spigelman AD, Williams CB, Talbot IC, et al. Upper gastrointestinal cancer in patients with familial adenomatous polyposis. *Lancet.* 1989; 2:783–5. [PubMed: 2571019]
155. Groves CJ, Saunders BP, Spigelman AD, et al. Duodenal cancer in patients with familial adenomatous polyposis (FAP): results of a 10 year prospective study. *Gut.* 2002; 50:636–41. [PubMed: 11950808]
156. Saurin JC, Gutknecht C, Napoleon B, et al. Surveillance of duodenal adenomas in familial adenomatous polyposis reveals high cumulative risk of advanced disease. *J Clin Oncol.* 2004; 22:493–8. [PubMed: 14752072]
157. Burt RW. Gastric fundic gland polyps. *Gastroenterology.* 2003; 125:1462–9. [PubMed: 14598262]
158. Weston BR, Helper DJ, Rex DK. Positive predictive value of endoscopic features deemed typical of gastric fundic gland polyps. *J Clin Gastroenterol.* 2003; 36:399–402. [PubMed: 12702980]
159. Bianchi LK, Burke CA, Bennett AE, et al. Fundic gland polyp dysplasia is common in familial adenomatous polyposis. *Clin Gastroenterol Hepatol.* 2008; 6:180–5. [PubMed: 18237868]
160. Zwick A, Munir M, Ryan CK, et al. Gastric adenocarcinoma and dysplasia in fundic gland polyps of a patient with attenuated adenomatous polyposis coli. *Gastroenterology.* 1997; 113:659–63. [PubMed: 9247488]
161. Hofgartner WT, Thorp M, Ramus MW, et al. Gastric adenocarcinoma associated with fundic gland polyps in a patient with attenuated familial adenomatous polyposis. *Am J Gastroenterol.* 1999; 94:2275–81. [PubMed: 10445562]
162. Attard TM, Giardiello FM, Argani P, et al. Fundic gland polyposis with high-grade dysplasia in a child with attenuated familial adenomatous polyposis and familial gastric cancer. *J Pediatr Gastroenterol Nutr.* 2001; 32:215–8. [PubMed: 11321399]
163. Sawada T, Muto T. Role of upper gastrointestinal surveillance in patients with familial adenomatous polyposis. *Gastrointest Endosc Clin N Am.* 1997; 7:99–111. [PubMed: 8995116]
164. Wallace MH, Phillips RK. Upper gastrointestinal disease in patients with familial adenomatous polyposis. *Br J Surg.* 1998; 85:742–50. [PubMed: 9667698]
165. Kadmon M, Tandara A, Herfarth C. Duodenal adenomatosis in familial adenomatous polyposis coli. A review of the literature and results from the Heidelberg Polyposis Register. *Int J Colorectal Dis.* 2001; 16:63–75. [PubMed: 11355321]
166. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®). *Colorectal Cancer Screening V.1.2010.* © National Comprehensive Cancer Network, Inc. 2014. All rights reserved. [cited 2014 March 13]; To view the most recent and complete version of the guideline, go online to NCCN.org. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, NCCN GUIDELINES®, and all other NCCN Content are trademarks owned by the National Comprehensive Cancer Network, Inc]. Available from http://www.nccn.org/professionals/physician_gls/f_guidelines.asp.
167. Kurtz RC, Sternberg SS, Miller HH, et al. Upper gastrointestinal neoplasia in familial polyposis. *Dig Dis Sci.* 1987; 32:459–65. [PubMed: 3032535]
168. Jagelman DG, DeCosse JJ, Bussey HJ. Upper gastrointestinal cancer in familial adenomatous polyposis. *Lancet.* 1988; 1:1149–51. [PubMed: 2896968]
169. Walsh N, Qizilbash A, Banerjee R, et al. Biliary neoplasia in Gardner's syndrome. *Arch Pathol Lab Med.* 1987; 111:76–7. [PubMed: 3800608]
170. Brevet M, Brehant O, Dumont F, et al. [Adenomatous polyposis of the gallbladder and Gardner's syndrome. A rare association]. *Gastroenterol Clin Biol.* 2007; 31:425–7. [PubMed: 17483782]

171. Herraiz M, Barbesino G, Faquin W, et al. Prevalence of thyroid cancer in familial adenomatous polyposis syndrome and the role of screening ultrasound examinations. *Clin Gastroenterol Hepatol.* 2007; 5:367–73. [PubMed: 17258512]
172. Cetta F, Montalto G, Gori M, et al. Germline mutations of the APC gene in patients with familial adenomatous polyposis-associated thyroid carcinoma: results from a European cooperative study. *J Clin Endocrinol Metabol.* 2000; 85:286–92.
173. Giardiello FM, Petersen GM, Brensinger JD, et al. Hepatoblastoma and APC gene mutation in familial adenomatous polyposis. *Gut.* 1996; 39:867–9. [PubMed: 9038672]
174. Vogt S, Jones N, Christian D, et al. Expanded extracolonic tumor spectrum in MUTYH-associated polyposis. *Gastroenterology.* 2009; 137:1976–85. [PubMed: 19732775]
175. Parc YR, Olschwang S, Desaint B, et al. Familial adenomatous polyposis: prevalence of adenomas in the ileal pouch after restorative proctocolectomy. *Ann Surg.* 2001; 233:360–4. [PubMed: 11224623]
176. Groves CJ, Beveridge G, Swain DJ, et al. Prevalence and morphology of pouch and ileal adenomas in familial adenomatous polyposis. *Dis Colon Rectum.* 2005; 48:816–23. [PubMed: 15747076]
177. Friederich P, de Jong AE, Mathus-Vliegen LM, et al. Risk of developing adenomas and carcinomas in the ileal pouch in patients with familial adenomatous polyposis. *Clin Gastroenterol Hepatol.* 2008; 6:1237–42. [PubMed: 18848811]
178. Tajika M, Nakamura T, Nakahara O, et al. Prevalence of adenomas and carcinomas in the ileal pouch after proctocolectomy in patients with familial adenomatous polyposis. *J Gastrointest Surg.* 2009; 13:1266–73. [PubMed: 19333660]
179. Campos FG, Imperiale AR, Seid VE, et al. Rectal and pouch recurrences after surgical treatment for familial adenomatous polyposis. *J Gastrointest Surg.* 2009; 13:129–36. [PubMed: 18766422]
180. Church J. Ileoanal pouch neoplasia in familial adenomatous polyposis: an underestimated threat. *Dis Colon Rectum.* 2005; 48:1708–13. [PubMed: 15937627]
181. Jeghers H, McKusick VA, Katz KH. Generalized intestinal polyposis and melanin spots of the oral mucosa, lips and digits. A syndrome of diagnostic significance. *N Engl J Med.* 1949; 241:993. [PubMed: 15399020]
182. Beggs AD, Latchford AR, Vasen HF, et al. Peutz-Jeghers syndrome: a systematic review and recommendations for management. *Gut.* 2010; 59:975–86. [PubMed: 20581245]
183. Schreiber IR, Baker M, Amos C, et al. The hamartomatous polyposis syndromes: a clinical and molecular review. *Am J Gastroenterol.* 2005; 100:476–90. [PubMed: 15667510]
184. Jaspersion K, Kohlmann W, Burt R. Inherited risk for colorectal cancer: practical approaches for identification, referral and management. *Pract Gastroenterol.* 2007; 31:37.
185. Giardiello FM, Trimbath JD. Peutz-Jeghers syndrome and management recommendations. *Clin Gastroenterol Hepatol.* 2006; 4:408–15. [PubMed: 16616343]
186. McGarrity TJ, Kulin HE, Zaino RJ. Peutz-Jeghers syndrome. *Am J Gastroenterol.* 2000; 95:596–604. [PubMed: 10710046]
187. Amos CI, Keitheri-Cheteri MB, Sabripour M, et al. Genotype-phenotype correlations in Peutz-Jeghers syndrome. *J Hum Genet.* 2004; 41:327–33.
188. van Lier MG, Mathus-Vliegen EM, Wagner A, et al. High cumulative risk of intussusception in patients with peutz-jeghers syndrome: time to update surveillance guidelines? *Am J Gastroenterol.* 2011; 106:940–5. [PubMed: 21157440]
189. Aretz S, Stienen D, Uhlhaas S, et al. High proportion of large genomic STK11 deletions in Peutz-Jeghers syndrome. *Hum Mutat.* 2005; 26:513–9. [PubMed: 16287113]
190. Volikos E, Robinson J, Aittomaki K, et al. LKB1 exonic and whole gene deletions are a common cause of Peutz-Jeghers syndrome. *J Med Genet.* 2006; 43:e18. [PubMed: 16648371]
191. Mehenni H, Resta N, Guanti G, et al. Molecular and clinical characteristics in 46 families affected with Peutz-Jeghers syndrome. *Dig Dis Sci.* 2007; 52:1924–33. [PubMed: 17404884]
192. Calva D, Howe JR. Hamartomatous polyposis syndromes. *Surg Clin North Am.* 2008; 88:779–817. vii. [PubMed: 18672141]

193. Gammon A, Jasperson K, Kohlmann W, et al. Hamartomatous polyposis syndromes. *Best Pract Res Clin Gastroenterol*. 2009; 23:219–31. [PubMed: 19414148]
194. Hearle N, Schumacher V, Menko FH, et al. Frequency and spectrum of cancers in the Peutz-Jeghers syndrome. *Clin Cancer Res*. 2006; 12:3209–15. [PubMed: 16707622]
195. Mehenni H, Resta N, Park JG, et al. Cancer risks in LKB1 germline mutation carriers. *Gut*. 2006; 55:984–90. [PubMed: 16407375]
196. van Lier MG, Wagner A, Mathus-Vliegen EM, et al. High cancer risk in Peutz-Jeghers syndrome: a systematic review and surveillance recommendations. *Am J Gastroenterol*. 2010; 105:1258–64. author reply 1265. [PubMed: 20051941]
197. Giardiello FM, Brensinger JD, Tersmette AC, et al. Very high risk of cancer in familial Peutz-Jeghers syndrome. *Gastroenterology*. 2000; 119:1447–53. [PubMed: 11113065]
198. Lim W, Olschwang S, Keller JJ, et al. Relative frequency and morphology of cancers in *stk11* mutation carriers. *Gastroenterology*. 2004; 126:1788–94. [PubMed: 15188174]
199. Ulbright TM, Amin MB, Young RH. Intratubular large cell hyalinizing sertoli cell neoplasia of the testis: a report of 8 cases of a distinctive lesion of the Peutz-Jeghers syndrome. *Am J Surg Pathol*. 2007; 31:827–35. [PubMed: 17527069]
200. You YN, Wolff BG, Boardman LA, et al. Peutz-Jeghers syndrome: a study of long-term surgical morbidity and causes of mortality. *Fam Cancer*. 2010; 9:609–16. [PubMed: 20571886]
201. Amaro R, Diaz G, Schneider J, et al. Peutz-Jeghers syndrome managed with a complete intraoperative endoscopy and extensive polypectomy. *Gastrointest Endosc*. 2000; 52:552–4. [PubMed: 11023581]
202. Parsi MA, Burke CA. Utility of capsule endoscopy in Peutz-Jeghers syndrome. *Gastrointest Endosc Clin N Am*. 2004; 14:159–67. [PubMed: 15062389]
203. Burke CA, Santisi J, Church J, et al. The utility of capsule endoscopy small bowel surveillance in patients with polyposis. *Am J Gastroenterol*. 2005; 100:1498–502. [PubMed: 15984971]
204. Ohmiya N, Taguchi A, Shirai K, et al. Endoscopic resection of Peutz-Jeghers polyps throughout the small intestine at double-balloon enteroscopy without laparotomy. *Gastrointest Endosc*. 2005; 61:140–7. [PubMed: 15672077]
205. May A, Nachbar L, Ell C. Double-balloon enteroscopy (push-and-pull enteroscopy) of the small bowel: feasibility and diagnostic and therapeutic yield in patients with suspected small bowel disease. *Gastrointest Endosc*. 2005; 62:62–70. [PubMed: 15990821]
206. Udd L, Katajisto P, Rossi DJ, et al. Suppression of Peutz-Jeghers polyposis by inhibition of cyclooxygenase-2. *Gastroenterology*. 2004; 127:1030–7. [PubMed: 15480979]
207. Wei C, Amos CI, Zhang N, et al. Suppression of Peutz-Jeghers polyposis by targeting mammalian target of rapamycin signaling. *Clin Cancer Res*. 2008; 14:1167–71. [PubMed: 18281551]
208. Chow E, Macrae F. A review of juvenile polyposis syndrome. *J Gastroenterol Hepatol*. 2005; 20:1634–40. [PubMed: 16246179]
209. Boardman LA. Heritable colorectal cancer syndromes: recognition and preventive management. *Gastroenterol Clin North Am*. 2002; 31:1107–31. [PubMed: 12489281]
210. Latchford AR, Neale K, Phillips RK, et al. Juvenile polyposis syndrome: a study of genotype, phenotype, and long-term outcome. *Dis Colon Rectum*. 2012; 55:1038–43. [PubMed: 22965402]
211. Howe JR, Bair JL, Sayed MG, et al. Germline mutations of the gene encoding bone morphogenetic protein receptor 1A in juvenile polyposis. *Nat Genet*. 2001; 28:184–7. [PubMed: 11381269]
212. Howe JR, Ringold JC, Summers RW, et al. A gene for familial juvenile polyposis maps to chromosome 18q21.1. *Am J Hum Genet*. 1998; 62:1129–36. [PubMed: 9545410]
213. Howe JR, Roth S, Ringold JC, et al. Mutations in the SMAD4/DPC4 gene in juvenile polyposis. *Science*. 1998; 280:1086–8. [PubMed: 9582123]
214. Zhou XP, Woodford-Richens K, Lehtonen R, et al. Germline mutations in BMPRIA/ALK3 cause a subset of cases of juvenile polyposis syndrome and of Cowden and Bannayan-Riley-Ruvalcaba syndromes. *Am J Hum Genet*. 2001; 69:704–11. [PubMed: 11536076]

215. Aretz S, Stienen D, Uhlhaas S, et al. High proportion of large genomic deletions and a genotype phenotype update in 80 unrelated families with juvenile polyposis syndrome. *J Med Genet.* 2007; 44:702–9. [PubMed: 17873119]
216. Calva-Cerqueira D, Dahdaleh FS, Woodfield G, et al. Discovery of the BMPRI1A promoter and germline mutations that cause juvenile polyposis. *Hum Mol Genet.* 2010; 19:4654–62. [PubMed: 20843829]
217. van Hattem WA, Brosens LA, de Leng WW, et al. Large genomic deletions of SMAD4, BMPRI1A and PTEN in juvenile polyposis. *Gut.* 2008; 57:623–7. [PubMed: 18178612]
218. He XC, Zhang J, Tong WG, et al. BMP signaling inhibits intestinal stem cell self-renewal through suppression of Wnt-beta-catenin signaling. *Nat Genet.* 2004; 36:1117–21. [PubMed: 15378062]
219. Hardwick JC, Van Den Brink GR, Bleuming SA, et al. Bone morphogenetic protein 2 is expressed by, and acts upon, mature epithelial cells in the colon. *Gastroenterology.* 2004; 126:111–21. [PubMed: 14699493]
220. Brosens LA, van Hattem A, Hyland LM, et al. Risk of colorectal cancer in juvenile polyposis. *Gut.* 2007; 56:965–7. [PubMed: 17303595]
221. Aaltonen, LA.; Jass, JR.; Howe, JR. Juvenile polyposis.. In: Hamilton, SR.; Aaltonen, LA., editors. *World Health Organization Classification of Tumours. Pathology and Genetics of Tumours of the Digestive System* IARC Press; Lyon: 2000. p. 130-2.
222. Wirtzfeld DA, Petrelli NJ, Rodriguez-Bigas MA. Hamartomatous polyposis syndromes: molecular genetics, neoplastic risk, and surveillance recommendations. *Ann Surg Oncol.* 2001; 8:319–27. [PubMed: 11352305]
223. Oncel M, Church JM, Remzi FH, et al. Colonic surgery in patients with juvenile polyposis syndrome: a case series. *Dis Colon Rectum.* 2005; 48:49–55. discussion 55-6. [PubMed: 15690657]
224. Heald B, Mester J, Rybicki L, et al. Frequent gastrointestinal polyps and colorectal adenocarcinomas in a prospective series of PTEN mutation carriers. *Gastroenterology.* 2010; 139:1927–33. [PubMed: 20600018]
225. Stanich PP, Owens VL, Sweetser S, et al. Colonic polyposis and neoplasia in Cowden syndrome. *Mayo Clin Proc.* 2011; 86:489–92. [PubMed: 21628613]
226. Trufant JW, Greene L, Cook DL, et al. Colonic ganglioneuromatous polyposis and metastatic adenocarcinoma in the setting of Cowden syndrome: a case report and literature review. *Hum Pathol.* 2012; 43:601–4. [PubMed: 22018627]
227. Levi Z, Baris HN, Kedar I, et al. Upper and lower gastrointestinal findings in PTEN mutation-positive Cowden syndrome patients participating in an active surveillance program. *Clin Transl Gastroenterol.* 2011; 17:4.
228. Pilarski R. Cowden syndrome: a critical review of the clinical literature. *J Genet Couns.* 2009; 18:13–27. [PubMed: 18972196]
229. McGarrity TJ, Wagner Baker MJ, Ruggiero FM, et al. GI polyposis and glycogenic acanthosis of the esophagus associated with PTEN mutation positive Cowden syndrome in the absence of cutaneous manifestations. *Am J Gastroenterol.* 2003; 98:1429–34. [PubMed: 12818292]
230. Al-Th ihli K, Palma L, Marcus V, et al. A case of Cowden's syndrome presenting with gastric carcinomas and gastrointestinal polyposis. *Nat Clin Pract Gastroenterol Hepatol.* 2009; 6:184–9. [PubMed: 19190598]
231. Liaw D, Marsh DJ, Li J, et al. Germline mutations of the PTEN gene in Cowden disease, an inherited breast and thyroid cancer syndrome. *Nat Genet.* 1997; 16:64–7. [PubMed: 9140396]
232. Lynch ED, Ostermeyer EA, Lee MK, et al. Inherited mutations in PTEN that are associated with breast cancer, Cowden disease, and juvenile polyposis. *Am J Hum Genet.* 1997; 61:1254–60. [PubMed: 9399897]
233. Nelen MR, Padberg GW, Peeters EA, et al. Localization of the gene for Cowden disease to chromosome 10q22-23. *Nat Genet.* 1996; 13:114–6. [PubMed: 8673088]
234. Nelen MR, van Staveren WC, Peeters EA, et al. Germline mutations in the PTEN/MMAC1 gene in patients with Cowden disease. *Hum Mol Genet.* 1997; 6:1383–7. [PubMed: 9259288]
235. Tan MH, Mester JL, Ngeow J, et al. Lifetime cancer risks in individuals with germline PTEN mutations. *Clin Cancer Res.* 2012; 18:400–7. [PubMed: 22252256]

236. Riegert-Johnson DL, Gleeson FC, Roberts M, et al. Cancer and Lhermitte-Duclos disease are common in Cowden syndrome patients. *Hered Cancer Clin Pract*. 2010; 8:1897–4287.
237. Hyman NH, Anderson P, Blasyk H. Hyperplastic polyposis and the risk of colorectal cancer. *Dis Colon Rectum*. 2004; 47:2101–4. [PubMed: 15657661]
238. Leggett BA, Devereaux B, Biden K, et al. Hyperplastic polyposis: association with colorectal cancer. *Am J Surg Pathol*. 2001; 25:177–84. [PubMed: 11176066]
239. Rex DK, Ahnen DJ, Baron JA, et al. Serrated lesions of the colorectum: review and recommendations from an expert panel. *Am J Gastroenterol*. 2012; 107:1315–29. quiz 1314, 1330. [PubMed: 22710576]
240. Burt, R.; Jass, JR. Hyperplastic polyposis.. In: Hamilton, SR.; Aaltonen, LA., editors. *World Health Organization Classification of Tumours. Pathology and Genetics of Tumours of the Digestive System* IARC Press; Lyon: 2000. p. 135-6.
241. Chow E, Lipton L, Lynch E, et al. Hyperplastic polyposis syndrome: phenotypic presentations and the role of MBD4 and MYH. *Gastroenterology*. 2006; 131:30–9. [PubMed: 16831587]
242. Snover, DC.; Ahnen, DJ.; Burt, RW., et al. Serrated polyps of the colon and rectum and serrated polyposis.. In: Bosman, FT.; Carneiro, F.; Hruban, RH., et al., editors. *WHO Classification of Tumours of the Digestive System* International Agency for Research on Cancer (IARC). Lyon: 2010. p. 160-5.
243. Orłowska J. Hyperplastic polyposis syndrome and the risk of colorectal cancer. *Gut*. 2012; 61:470–1. [PubMed: 21636642]
244. Biswas S, Ellis AJ, Guy R, et al. High prevalence of hyperplastic polyposis syndrome (serrated polyposis) in the NHS bowel cancer screening programme. *Gut*. 2013; 62:475. [PubMed: 22851664]
245. Moreira L, Pellisé M, Carballal S, et al. High prevalence of serrated polyposis syndrome in FIT-based colorectal cancer screening programmes. *Gut*. 2013; 62:476–7. [PubMed: 22982987]
246. Buchanan D, Young J. A perspective on bi-allelic MUTYH mutations in patients with hyperplastic polyposis syndrome. *Gastroenterology*. 2009; 136:2407–8. 2009. [PubMed: 19406141]
247. Walker RG, Landmann JK, Hewett DG, et al. Hyperplastic polyposis syndrome is associated with cigarette smoking, which may be a modifiable risk factor. *Am J Gastroenterol*. 2010; 105:1642–7. [PubMed: 20125129]
248. Buchanan DD, Sweet K, Drini M, et al. Phenotypic diversity in patients with multiple serrated polyps: a genetics clinic study. *Int J Colorectal Dis*. 2010; 25:703–12. [PubMed: 20213458]
249. Kalady MF, Jarrar A, Leach B, et al. Defining phenotypes and cancer risk in hyperplastic polyposis syndrome. *Dis Colon Rectum*. 2011; 54:164–70. [PubMed: 21228663]
250. Win AK, Walters RJ, Buchanan DD, et al. Cancer risks for relatives of patients with serrated polyposis. *Am J Gastroenterol*. 2012; 107:770–8. [PubMed: 22525305]
251. Lanspa SJ, Ahnen DJ, Lynch HT. Editorial: serrated polyposis: the last (or only the latest?) frontier of familial polyposis? *Am J Gastroenterol*. 2012; 107:779–81. [PubMed: 22552244]
252. Rosty C, Parry S, Young JP. Serrated polyposis: an enigmatic model of colorectal cancer predisposition. *Patholog Res Int*. 2011; 2011:157073. [PubMed: 21660283]
253. Boparai KS, Mathus-Vliegen EM, Koornstra JJ, et al. Increased colorectal cancer risk during follow-up in patients with hyperplastic polyposis syndrome: a multicentre cohort study. *Gut*. 2009; 59:1094–100. [PubMed: 19710031]
254. Rosty C, Buchanan DD, Walsh MD, et al. Phenotype and polyp landscape in serrated polyposis syndrome: a series of 100 patients from genetics clinics. *Am J Surg Pathol*. 2012; 36:876–82. [PubMed: 22510757]
255. Rosty C, Walsh MD, Walters RJ, et al. Multiplicity and molecular heterogeneity of colorectal carcinomas in individuals with serrated polyposis. *Am J Surg Pathol*. 2013; 37:434–42. 10.1097/PAS.0b013e318270f748. [PubMed: 23211288]
256. Young JP, Parry S. Risk factors: hyperplastic polyposis syndrome and risk of colorectal cancer. *Nat Rev Gastroenterol Hepatol*. 2010; 7:594–5. [PubMed: 21042338]

257. Hazewinkel Y, Reitsma JB, Nagengast FM, et al. Extracolonic cancer risk in patients with serrated polyposis syndrome and their first-degree relatives. *Fam Cancer*. 2013; 12:669–73. [PubMed: 23591707]
258. Oquinená S, Guerra A, Pueyo A, et al. Serrated polyposis: prospective study of first-degree relatives. *Eur J Gastroenterol Hepatol*. 2013; 25:28–32. [PubMed: 23011040]
259. Boparai KS, Reitsma JB, Lemmens V, et al. Increased colorectal cancer risk in first-degree relatives of patients with hyperplastic polyposis syndrome. *Gut*. 2010; 59:1222–5. [PubMed: 20584785]
260. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Genetic/Familial High Risk Assessment: Colorectal V.1.2013. © National Comprehensive Cancer Network, Inc. 2014. All rights reserved. [cited 2014 March 13]; To view the most recent and complete version of the guideline, go online to NCCN.org. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, NCCN GUIDELINES®, and all other NCCN Content are trademarks owned by the National Comprehensive Cancer Network, Inc]. Available from http://www.nccn.org/professionals/physician_gls/f_guidelines.asp.
261. Leachman SA, Carucci J, Kohlmann W, et al. Selection criteria for genetic assessment of patients with familial melanoma. *J Am Acad Dermatol*. 2009; 61:677–e1-677. e14. [PubMed: 19751883]
262. Brand RE, Lerch MM, Rubinstein WS, et al. Advances in counselling and surveillance of patients at risk for pancreatic cancer. *Gut*. 2007; 56:1460–9. [PubMed: 17872573]
263. Ellis I, Lerch MM, Whitcomb DC. Genetic testing for hereditary pancreatitis: guidelines for indications, counselling, consent and privacy issues. *Pancreatology*. 2001; 1:405–15. [PubMed: 12120217]
264. Lowenfels AB, Maisonneuve P, DiMagno EP, et al. Hereditary pancreatitis and the risk of pancreatic cancer. International Hereditary Pancreatitis Study Group. *J Natl Cancer Inst*. 1997; 89:442–6. [PubMed: 9091646]
265. Vasen HF, Gruis NA, Frants RR, et al. Risk of developing pancreatic cancer in families with familial atypical multiple mole melanoma associated with a specific 19 deletion of p16 (p16-Leiden). *Int J Cancer*. 2000; 87:809–11. [PubMed: 10956390]
266. Goldstein AM, Fraser MC, Struwing JP, et al. Increased risk of pancreatic cancer in melanoma-prone kindreds with p16INK4 mutations. *N Engl J Med*. 1995; 333:970–4. [PubMed: 7666916]
267. Borg A, Sandberg T, Nilsson K, et al. High frequency of multiple melanomas and breast and pancreas carcinomas in CDKN2A mutation-positive melanoma families. *J Natl Cancer Inst*. 2000; 92:1260–6. [PubMed: 10922411]
268. Thompson D, Easton DF. Breast Cancer Linkage Consortium. Cancer incidence in BRCA1 mutation carriers. *J Natl Cancer Inst*. 2002; 94:1358–65. [PubMed: 12237281]
269. BCLC. The Breast Cancer Linkage Consortium. Cancer risks in BRCA2 mutation carriers. *J Natl Cancer Inst*. 1999; 91:1310–6. [PubMed: 10433620]
270. van Asperen CJ, Brohet RM, Meijers-Heijboer EJ, et al. Cancer risks in BRCA2 families: estimates for sites other than breast and ovary. *J Med Genet*. 2005; 42:711–9. [PubMed: 16141007]
271. Kastrinos F, Mukherjee B, Tayob N, et al. Risk of pancreatic cancer in families with Lynch syndrome. *JAMA*. 2009; 302:1790–5. [PubMed: 19861671]
272. Geoffroy-Perez B, Janin N, Ossian K, et al. Cancer risk in heterozygotes for ataxia-telangiectasia. *Int J Cancer*. 2001; 93:288–93. [PubMed: 11410879]
273. Brand RE, Lynch HT. Hereditary pancreatic adenocarcinoma: a clinical perspective. *Med Clin N Am*. 2000; 84:665–75. [PubMed: 10872423]
274. Permuth-Wey J, Egan KM. Family history is a significant risk factor for pancreatic cancer: results from a systematic review and meta-analysis. *Fam Cancer*. 2009; 8:109–17. [PubMed: 18763055]
275. Klein AP, Beaty TH, Bailey-Wilson JE, et al. Evidence for a major gene influencing risk of pancreatic cancer. *Genet Epidemiol*. 2002; 23:133–49. [PubMed: 12214307]
276. Klein AP, Brune KA, Petersen GM, et al. Prospective risk of pancreatic cancer in familial pancreatic cancer kindreds. *Cancer Res*. 2004; 64:2634–8. [PubMed: 15059921]
277. Brune KA, Lau B, Palmisano E, et al. Importance of age of onset in pancreatic cancer kindreds. *J Natl Cancer Inst*. 2010; 102:119–26. [PubMed: 20068195]

278. Wang W, Chen S, Brune KA, et al. PancPRO: risk assessment for individuals with a family history of pancreatic cancer. *J Clin Oncol*. 2007; 25:1417–22. [PubMed: 17416862]
279. Lal G, Liu G, Schmocker B, et al. Inherited predisposition to pancreatic adenocarcinoma: role of family history and germ-line p16, BRCA1, and BRCA2 mutations. *Cancer Res*. 2000; 60:409–16. [PubMed: 10667595]
280. McWilliams RR, Wieben ED, Rabe KG, et al. Prevalence of CDKN2A mutations in pancreatic cancer patients: implications for genetic counseling. *Eur J Hum Genet*. 2011; 19:472–8. [PubMed: 21150883]
281. Ghiorzo P, Fornarini G, Sciallero S, et al. CDKN2A is the main susceptibility gene in Italian pancreatic cancer families. *J Med Genet*. 2012; 49:164–70. [PubMed: 22368299]
282. Pogue-Geile KL, Chen R, Bronner MP, et al. Palladin mutation causes familial pancreatic cancer and suggests a new cancer mechanism. *PLoS Med*. 2006; 3:e516. [PubMed: 17194196]
283. Roberts NJ, Jiao Y, Yu J, et al. ATM mutations in patients with hereditary pancreatic cancer. *Cancer Discov*. 2012; 2:41–6. [PubMed: 22585167]
284. Harinck F, Kluijdt I, van Mil SE, et al. Routine testing for PALB2 mutations in familial pancreatic cancer families and breast cancer families with pancreatic cancer is not indicated. *Eur J Hum Genet*. 2012; 20:577–9. [PubMed: 22166947]
285. Jones S, Hruban RH, Kamiyama M, et al. Exomic sequencing identifies PALB2 as a pancreatic cancer susceptibility gene. *Science*. 2009; 324:217. [PubMed: 19264984]
286. Axilbund JE, Argani P, Kamiyama M, et al. Absence of germline BRCA1 mutations in familial pancreatic cancer patients. *Cancer Biol Ther*. 2009; 8:131–5. [PubMed: 19029836]
287. Couch FJ, Johnson MR, Rabe KG, et al. The prevalence of BRCA2 mutations in familial pancreatic cancer. *Cancer Epidemiol Biomarkers Prev*. 2007; 16:342–6. [PubMed: 17301269]
288. Schneider R, Slater EP, Sina M, et al. German national case collection for familial pancreatic cancer (FaPaCa): ten years experience. *Fam Cancer*. 2011; 10:323–30. [PubMed: 21207249]
289. Klein AP, Borges M, Griffith M, et al. Absence of deleterious palladin mutations in patients with familial pancreatic cancer. *Cancer Epidemiol Biomarkers Prev*. 2009; 18:1328–30. [PubMed: 19336541]
290. Slater E, Amrillaeva V, Fendrich V, et al. Palladin mutation causes familial pancreatic cancer: absence in European families. *PLoS Med*. 2007; 4:e164. [PubMed: 17455999]
291. ACS. American Cancer Society. Cancer Facts & Figures 2013. 2013. [cited] Available from <http://www.cancer.org/acs/groups/content/@epidemiologysurveillance/documents/document/acspsc-036845.pdf>
292. Bartsch DK, Gress TM, Langer P. Familial pancreatic cancer—current knowledge. *Nat Rev Gastroenterol Hepatol*. 2012; 9:445–53. [PubMed: 22664588]
293. Grover S, Syngal S. Hereditary pancreatic cancer. *Gastroenterology*. 2010; 139:1076–80. [PubMed: 20727885]
294. Nishihara R, Wu K, Lochhead P, et al. Long-term colorectal-cancer incidence and mortality after lower endoscopy. *N Engl J Med*. 2013; 369:1095–105. [PubMed: 24047059]
295. Canto MI, Hruban RH, Fishman EK, et al. Frequent detection of pancreatic lesions in asymptomatic high-risk individuals. *Gastroenterology*. 2012; 142:796–804. quiz e14-5. [PubMed: 22245846]
296. Ludwig E, Olson SH, Bayuga S, et al. Feasibility and yield of screening in relatives from familial pancreatic cancer families. *Am J Gastroenterol*. 2011; 106:946–54. [PubMed: 21468009]
297. Anderson MA, Zolotarevsky E, Cooper KL, et al. Alcohol and tobacco lower the age of presentation in sporadic pancreatic cancer in a dose-dependent manner: a multicenter study. *Am J Gastroenterol*. 2012; 107:1730–9. [PubMed: 22929760]
298. Brand RE, Greer JB, Zolotarevsky E, et al. Pancreatic cancer patients who smoke and drink are diagnosed at younger ages. *Clin Gastroenterol Hepatol*. 2009; 7:1007–12. [PubMed: 19560558]
299. Lowenfels AB, Maisonneuve P, Whitcomb DC, et al. Cigarette smoking as a risk factor for pancreatic cancer in patients with hereditary pancreatitis. *JAMA*. 2001; 286:169–70. [PubMed: 11448279]

300. Vasen HF, Wasser M, van Mil A, et al. Magnetic resonance imaging surveillance detects early-stage pancreatic cancer in carriers of a p16-Leiden mutation. *Gastroenterology*. 2011; 140:850–6. [PubMed: 21129377]
301. Tanaka M, Fernandez-del Castillo C, Adsay V, et al. International consensus guidelines 2012 for the management of IPMN and MCN of the pancreas. *Pancreatology*. 2012; 12:183–97. [PubMed: 22687371]
302. Rulyak SJ, Lowenfels AB, Maisonneuve P, et al. Risk factors for the development of pancreatic cancer in familial pancreatic cancer kindreds. *Gastroenterology*. 2003; 124:1292–9. [PubMed: 12730869]
303. NCI. National Cancer Institute. SEER Cancer Statistics Factsheets: Stomach Cancer. 2013. [cited] Available from <http://seer.cancer.gov/statfacts/html/stomach.html>
304. Jones EG. Familial gastric cancer. *N Z Med J*. 1964; 63:287–96. [PubMed: 14158754]
305. Seevaratnam R, Coburn N, Cardoso R, et al. A systematic review of the indications for genetic testing and prophylactic gastrectomy among patients with hereditary diffuse gastric cancer. *Gastric Cancer*. 2012; 15:10.
306. Guilford P, Hopkins J, Harraway J, et al. E-cadherin germline mutations in familial gastric cancer. *Nature*. 1998; 392:402–5. [PubMed: 9537325]
307. Gayther SA, Goringe KL, Ramus SJ, et al. Identification of germ-line E-cadherin mutations in gastric cancer families of European origin. *Cancer Res*. 1998; 58:4086–9. [PubMed: 9751616]
308. Guilford PJ, Hopkins JB, Grady WM, et al. E-cadherin germline mutations define an inherited cancer syndrome dominated by diffuse gastric cancer. *Hum Mutat*. 1999; 14:249–55. [PubMed: 10477433]
309. Richards FM, McKee SA, Rajpar MH, et al. Germline E-cadherin gene (CDH1) mutations predispose to familial gastric cancer and colorectal cancer. *Hum Mol Genet*. 1999; 8:607–10. [PubMed: 10072428]
310. Blair V, Martin I, Shaw D, et al. Hereditary diffuse gastric cancer: diagnosis and management. *Clin Gastroenterol Hepatol*. 2006; 4:262–75. [PubMed: 16527687]
311. Pharoah PD, Guilford P, Caldas C. Incidence of gastric cancer and breast cancer in CDH1 (E-cadherin) mutation carriers from hereditary diffuse gastric cancer families. *Gastroenterology*. 2001; 121:1348–53. [PubMed: 11729114]
312. Fitzgerald RC, Hardwick R, Huntsman D, et al. Hereditary diffuse gastric cancer: updated consensus guidelines for clinical management and directions for future research. *J Med Genet*. 2010; 47:436–44. [PubMed: 20591882]
313. Brooks-Wilson AR, Kaurah P, Suriano G, et al. Germline E-cadherin mutations in hereditary diffuse gastric cancer: assessment of 42 new families and review of genetic screening criteria. *J Med Genet*. 2004; 41:508–17. [PubMed: 15235021]
314. Caldas C, Carneiro F, Lynch HT, et al. Familial gastric cancer: overview and guidelines for management. *J Med Genet*. 1999; 36:873–80. [PubMed: 10593993]
315. Kaurah, P.; Huntsman, DG. [21 June 2011] Hereditary diffuse gastric cancer.. *GeneReviews*. Nov 4. 2002 cited; available from <http://www.ncbi.nlm.nih.gov/books/NBK1139/>
316. Iwama T, Mishima Y, Utsunomiya J. The impact of familial adenomatous polyposis on the tumorigenesis and mortality at the several organs. Its rational treatment. *Ann Surg*. 1993; 217:101–8. [PubMed: 8382467]
317. Lynch HT, Smyrk T, McGinn T, et al. Attenuated familial adenomatous polyposis (AFAP). A phenotypically and genotypically distinctive variant of FAP. *Cancer*. 1995; 76:2427–33. [PubMed: 8625067]
318. Neklason DW, Stevens J, Boucher KM, et al. American founder mutation for attenuated familial adenomatous polyposis. *Clin Gastroenterol Hepatol*. 2008; 6:46–52. [PubMed: 18063416]
319. Nielsen M, Hes FJ, Nagengast FM, et al. Germline mutations in APC and MUTYH are responsible for the majority of families with attenuated familial adenomatous polyposis. *Clin Genet*. 2007; 71:427–33. [PubMed: 17489848]
320. Giardiello FM, Hamilton SR, Kern SE, et al. Colorectal neoplasia in juvenile polyposis or juvenile polyps. *Arch Dis Child*. 1991; 66:971–5. [PubMed: 1656892]

321. Howe JR, Mitros FA, Summers RW. The risk of gastrointestinal carcinoma in familial juvenile polyposis. *Ann Surg Oncol*. 1998; 5:751–6. [PubMed: 9869523]
322. Jass, J. Pathology of polyposis syndromes with special reference to juvenile polyposis.. In: Utsunomiya, J.; Lynch, H., editors. *Hereditary Colorectal Cancer*. Springer; Japan: 1990. p. 343-50.
323. Jass JR, Williams CB, Bussey HJ, et al. Juvenile polyposis--a precancerous condition. *Histopathology*. 1988; 13:619–30. [PubMed: 2853131]
324. Marra G, Armelao F, Vecchio FM, et al. Cowden's disease with extensive gastrointestinal polyposis. *J Clin Gastroenterol*. 1994; 18:42–7. [PubMed: 8113585]
325. Vasen HF, Griffioen G, Offerhaus GJ, et al. The value of screening and central registration of families with familial adenomatous polyposis. A study of 82 families in The Netherlands. *Dis Colon Rectum*. 1990; 33:227–30. [PubMed: 2155763]
326. Nascimbeni R, Pucciarelli S, Di Lorenzo D, et al. Rectum-sparing surgery may be appropriate for biallelic MUTYH-associated polyposis. *Dis Colon Rectum*. 2010; 53:1670–5. [PubMed: 21178863]
327. Nielsen M, van Steenberg LN, Jones N, et al. Survival of MUTYH-associated polyposis patients with colorectal cancer and matched control colorectal cancer patients. *J Natl Cancer Inst*. 2010; 102:1724–30. [PubMed: 21044966]
328. NCI. National Cancer Institute. SEER Cancer Statistics Review 1975-2010. 2013. [cited; Lifetime risk (percent) of being diagnosed with cancer by site]. Available from http://seer.cancer.gov/csr/1975_2010/results_merged/topic_lifetime_risk_diagnosis.pdf
329. Ponti G, Losi L, Pedroni M, et al. Value of MLH1 and MSH2 mutations in the appearance of Muir–Torre syndrome phenotype in HNPCC patients presenting sebaceous gland tumors or keratoacanthomas. *J Invest Dermatol*. 2006; 126:2302–7. [PubMed: 16826164]
330. Schwartz RA, Torre DP. The Muir-Torre syndrome: a 25-year retrospect. *J Am Acad Dermatol*. 1995; 33:90–104. [PubMed: 7601953]
331. South CD, Hampel H, Comeras I, et al. The frequency of Muir-Torre syndrome among Lynch syndrome families. *J Natl Cancer Inst*. 2008; 100:277–81. [PubMed: 18270343]
332. Grindedal EM, Møller P, Eeles R, et al. Germ-line mutations in mismatch repair genes associated with prostate cancer. *Cancer Epidemiol Biomark Prev*. 2009; 18:2460–7.
333. Heiskanen I, Kellokumpu I, Jarvinen H. Management of duodenal adenomas in 98 patients with familial adenomatous polyposis. *Endoscopy*. 1999; 31:412–6. [PubMed: 10494676]
334. Giardiello FM, Offerhaus GJ, Lee DH, et al. Increased risk of thyroid and pancreatic carcinoma in familial adenomatous polyposis. *Gut*. 1993; 34:1394–6. [PubMed: 8244108]
335. Truta B, Allen BA, Conrad PG, et al. Genotype and phenotype of patients with both familial adenomatous polyposis and thyroid carcinoma. *Fam Cancer*. 2003; 2:95–9. [PubMed: 14574158]
336. Hamilton SR, Liu B, Parsons RE, et al. The molecular basis of Turcot's syndrome. *N Engl J Med*. 1995; 332:839–47. [PubMed: 7661930]
337. Paraf F, Jothy S, Van Meir EG. Brain tumor-polyposis syndrome: two genetic diseases? *J Clin Oncol*. 1997; 15:2744–58. [PubMed: 9215849]
338. Van Meir EG. “Turcot's syndrome”: phenotype of brain tumors, survival and mode of inheritance. *Int J Cancer*. 1998; 75:162–4. [PubMed: 9426707] Korse SE, Harinck F, van Lier MG, et al. Pancreatic cancer risk in Peutz-Jeghers syndrome patients: a large cohort study and implications for surveillance. *J Med Genet*. 2013; 50:59–64. [PubMed: 23240097]
339. Chen KT. Female genital tract tumors in Peutz-Jeghers syndrome. *Hum Pathol*. 1986; 17:858–61. [PubMed: 3733073]
340. Brownstein MH, Wolf M, Bikowski JB. Cowden's disease: a cutaneous marker of breast cancer. *Cancer*. 1978; 41:2393–8. [PubMed: 657103]
341. Bubien V, Bonnet F, Brouste V, et al. High cumulative risks of cancer in patients with PTEN hamartoma tumour syndrome. *J Med Genet*. 2013; 50:255–63. [PubMed: 23335809]
342. Starink TM, van der Veen JP, Arwert F, et al. The Cowden syndrome: a clinical and genetic study in 21 patients. *Clin Genet*. 1986; 29:222–33. [PubMed: 3698331]
343. Eng C. Cowden syndrome. *J Genet Counsel*. 1997; 6:181–92.

344. Milas M, Mester J, Metzger R, et al. Should patients with Cowden syndrome undergo prophylactic thyroidectomy? *Surgery*. 2012; 152:1201–10. [PubMed: 23158187]
345. Eng C. Will the real Cowden syndrome please stand up: revised diagnostic criteria. *J Med Genet*. 2000; 37:828–30. [PubMed: 11073535]
346. Pilarski R, Eng C. Will the real Cowden syndrome please stand up (again)? Expanding mutational and clinical spectra of the PTEN hamartoma tumour syndrome. *J Med Genet*. 2004; 41:323–6. [PubMed: 15121767]
347. Rijcken FE, Mourits MJ, Kleibeuker JH, et al. Gynecologic screening in hereditary nonpolyposis colorectal cancer. *Gynecol Oncol*. 2003; 91:74–80. [PubMed: 14529665]
348. Lecuru F, Le Frere Belda MA, Bats AS, et al. Performance of office hysteroscopy and endometrial biopsy for detecting endometrial disease in women at risk of human non-polyposis colon cancer: a prospective study. *Int J Gynecol Cancer*. 2008; 18:1326–31. [PubMed: 18217965]
349. Gerritzen LH, Hoogerbrugge N, Oei AL, et al. Improvement of endometrial biopsy over transvaginal ultrasound alone for endometrial surveillance in women with Lynch syndrome. *Fam Cancer*. 2009; 8:391–7. [PubMed: 19504173]
350. Stuckless S, Green J, Dawson L, et al. Impact of gynecological screening in Lynch syndrome carriers with an MSH2 mutation. *Clin Genet*. 2013; 83:359–64. [PubMed: 22775459]
351. Renkonen-Sinisalo L, Sipponen P, Aarnio M, et al. No support for endoscopic surveillance for gastric cancer in hereditary non-polyposis colorectal cancer. *Scand J Gastroenterol*. 2002; 37:574–7. [PubMed: 12059060]
352. Biasco G, Nobili E, Calabrese C, et al. Impact of surgery on the development of duodenal cancer in patients with familial adenomatous polyposis. *Dis Colon Rectum*. 2006; 49:1860–6. [PubMed: 17103055]
353. Jarrar AM, Milas M, Mitchell J, et al. Screening for thyroid cancer in patients with familial adenomatous polyposis. *Ann Surg*. 2011; 253:515–21. [PubMed: 21173694]
354. Latchford AR, Neale K, Phillips RK, et al. Peutz-Jeghers syndrome: intriguing suggestion of gastrointestinal cancer prevention from surveillance. *Dis Colon Rectum*. 2011; 54:1547–51. [PubMed: 22067184]
355. Brown G, Fraser C, Schofield G, et al. Video capsule endoscopy in peutzjeghers syndrome: a blinded comparison with barium follow-through for detection of small-bowel polyps. *Endoscopy*. 2006; 38:385–90. [PubMed: 16680639]
356. Gupta A, Aftergate AJ, Burling D, et al. A prospective study of MR enterography versus capsule endoscopy for the surveillance of adult patients with Peutz-Jeghers syndrome. *AJR Am J Roentgenol*. 2010; 195:108–16. [PubMed: 20566803]
357. Poley JW, Kluijt I, Gouma DJ, et al. The yield of first-time endoscopic ultrasonography in screening individuals at a high risk of developing pancreatic cancer. *Am J Gastroenterol*. 2009; 104:2175–81. [PubMed: 19491823]
358. Lindor NM, Petersen GM, Hadley DW, et al. Recommendations for the care of individuals with an inherited predisposition to Lynch syndrome: a systematic review. *JAMA*. 2006; 296:1507–17. [PubMed: 17003399]
359. Lynch HT, Lynch JF, Lynch PM, et al. Hereditary colorectal cancer syndromes: molecular genetics, genetic counseling, diagnosis and management. *Fam Cancer*. 2008; 7:27–39. [PubMed: 17999161]
360. Faughnan ME, Palda VA, Garcia-Tsao G, et al. International guidelines for the diagnosis and management of hereditary haemorrhagic telangiectasia. *J Med Genet*. 2011; 48:73–87. [PubMed: 19553198]
361. Kastrinos F, Mukherjee B, Tayob N, et al. Risk of pancreatic cancer in families with Lynch syndrome. *JAMA*. 2009; 302:1790–5. [PubMed: 19861671]
362. Slater E, Amrillaeva V, Fendrich V, et al. Palladin mutation causes familial pancreatic cancer: absence in European families. *PLoS Med*. 2007; 4:e164. [PubMed: 17455999]
363. Robson ME, Storm CD, Weitzel J, et al. American Society of Clinical Oncology policy statement update: genetic and genomic testing for cancer susceptibility. *J Clin Oncol*. 2010; 28:893–901. [PubMed: 20065170]

Table 1

Summary statements

Standard for minimal cancer family history assessment in gastrointestinal (GI) practice

A family history of cancer and premalignant GI conditions that provides sufficient information to develop a preliminary determination of the risk of a familial predisposition to cancer should be obtained for all patients being evaluated in outpatient gastroenterology and endoscopy practices.

Essential elements of a family history include presence and type of cancer diagnoses in first- and second-degree relatives, and presence and (ideally) type of polyps in first-degree relatives; age and lineage should be noted for each diagnosis.

Lynch syndrome (LS)

All newly diagnosed colorectal cancers (CRCs) should be evaluated for mismatch repair deficiency.

Analysis may be done by immunohistochemical testing for the *MLH1/MSH2/MSH6/PMS2* proteins and/or testing for microsatellite instability. Tumors that demonstrate loss of *MLH1* should undergo BRAF testing or analysis for *MLH1* promoter hypermethylation.

Individuals who have a personal history of a tumor showing evidence of mismatch repair deficiency (and no demonstrated BRAF mutation or hypermethylation of *MLH1*), a known family mutation associated with LS, or a risk of ~5% chance of LS based on risk prediction models should undergo genetic evaluation for LS.

Genetic testing of patients with suspected LS should include germline mutation genetic testing for the *MLH1, MSH2, MSH6, PMS2*, and/or *EPCAM* genes or the altered gene(s) indicated by immunohistochemical (IHC) testing.

*Adenomatous polyposis syndromes**Familial adenomatous polyposis (FAP)/MUTYH-associated polyposis/attenuated polyposis*

Individuals who have a personal history of >10 cumulative colorectal adenomas, a family history of one of the adenomatous polyposis syndromes, or a history of adenomas and FAP-type extracolonic manifestations (duodenal/ampullary adenomas, desmoid tumors (abdominal>peripheral), papillary thyroid cancer, congenital hypertrophy of the retinal pigment epithelium ((CHRPE), epidermal cysts, osteomas) should undergo assessment for the adenomatous polyposis syndromes.

Genetic testing of patients with suspected adenomatous polyposis syndromes should include *APC* and *MUTYH* gene mutation analysis.

*Hamartomatous polyposis syndromes**Peutz–Jeghers syndrome (PJS)*

Individuals with perioral or buccal pigmentation and/or two or more histologically characteristic gastrointestinal hamartomatous polyp(s) or a family history of PJS should be evaluated for PJS.

Genetic evaluation of a patient with possible PJS should include testing for *STK11* mutations.

Juvenile polyposis syndrome (JPS)

Individuals with five or more juvenile polyps in the colorectum or any juvenile polyps in other parts of the GI tract should undergo evaluation for JPS.

Genetic evaluation of a patient with possible JPS should include testing for *SMAD4* and *BMPRIA* mutations.

Cowden syndrome (PTEN hamartoma tumor syndrome)

Individuals with multiple gastrointestinal hamartomas or ganglioneuromas should be evaluated for Cowden syndrome and related conditions.

Genetic evaluation of a patient with possible Cowden syndrome should include testing for *PTEN* mutations.

Serrated/hyperplastic polyposis syndrome

Individuals who meet at least one of the following criteria have the clinical diagnosis of serrated polyposis syndrome (SPS): (i) at least 5 serrated polyps proximal to the sigmoid colon with 2 of these being >10 mm; (ii) any number of serrated polyps proximal to the sigmoid colon in an individual who has a first-degree relative (FDR) with serrated polyposis; and (iii) >20 serrated polyps of any size, distributed throughout the large intestine.

A clear genetic etiology has not yet been defined for SPS, and therefore genetic testing is currently not routinely recommended for SPS patients; testing for *MUTYH* mutations may be considered for SPS patients with concurrent adenomas and/or a family history of adenomas.

Hereditary pancreatic cancer

Individuals should be considered to be at risk for familial pancreatic adenocarcinoma if they (i) have a known genetic syndrome associated with pancreatic cancer, including hereditary breast—ovarian cancer syndrome, familial atypical multiple melanoma and mole syndrome (FAMMM), PJS, LS, or other gene mutations associated with an increased risk of pancreatic adenocarcinoma; or (ii) have two relatives with pancreatic adenocarcinoma, where one is a FDR; (iii) have three or more relatives with pancreatic cancer; or (iv) have a history of hereditary pancreatitis.

Genetic testing of patients with suspected familial pancreatic cancer should include analysis of *BRCA1/2*, *CDKN2A*, *PALB2*, and *ATM*. Evaluation for PJS, LS, and hereditary pancreatitis-associated genes should be considered if other component personal and/or family history criteria are met for the syndrome.

Hereditary gastric cancer

Hereditary diffuse gastric cancer (HDGC)

Individuals with (i) 2 cases of diffuse gastric cancer, with at least one diagnosed at <50 years; (ii) 3 cases of documented diffuse cancer in first- or second degree relatives independent of age of onset; (iii) diffuse gastric cancer diagnosed at <40 years; (iv) a personal or family history of diffuse gastric cancer and lobular breast cancer with one diagnosed at <50 years should be evaluated for HDGC.

Genetic testing of individuals who fulfill HDGC clinical criteria should include analysis of *CDH1* mutations.

Table 2

Summary of recommendations

Lynch syndrome (LS)

1. In individuals at risk for or affected with LS, screening for colorectal cancer by colonoscopy should be performed at least every 2 years, beginning between ages 20 and 25 years. Annual colonoscopy should be considered in confirmed mutation carriers (strong recommendation, moderate quality of evidence for screening, and very low quality of evidence for annual surveillance and age of initiation).
2. Colectomy with ileorectal anastomosis (IRA) is the preferred treatment of patients affected with LS with colon cancer or colonic neoplasia not controllable by endoscopy. Segmental colectomy is an option in patients unsuitable for total colectomy if regular postoperative surveillance is conducted (conditional recommendation, moderate quality of evidence).
3. Hysterectomy and bilateral salpingo-oophorectomy should be offered to women who are known LS mutation carriers and who have finished child bearing, optimally at age 40–45 years (conditional recommendation, low quality of evidence).
4. Screening for endometrial cancer and ovarian cancer should be offered to women at risk for or affected with LS by endometrial biopsy and transvaginal ultrasound annually, starting at age 30 to 35 years before undergoing surgery or if surgery is deferred (conditional recommendation, very low quality of evidence).
5. Screening for gastric and duodenal cancer can be considered in individuals at risk for or affected with LS by baseline esophagogastroduodenoscopy (EGD) with gastric biopsy at age 30–35 years, and treatment of *H. pylori* infection when found. Data for ongoing regular surveillance are limited, but ongoing surveillance every 3–5 years may be considered if there is a family history of gastric or duodenal cancer (conditional recommendation, very low quality of evidence).
6. Screening beyond population-based recommendations for cancers of the urinary tract, pancreas, prostate, and breast is not recommended unless there is a family history of the specific cancers (conditional recommendation, low quality of evidence).
7. Although data suggest that daily aspirin may decrease the risk of colorectal and extracolonic cancer in LS, currently the evidence is not sufficiently robust or mature to make a recommendation for its standard use (conditional recommendation, moderate quality of evidence).

*Adenomatous polyposis syndromes**Familial adenomatous polyposis (FAP)/MUTYH-associated polyposis (MAP)/attenuated polyposis*

8. In individuals at risk for or affected with the classic AP syndromes, screening for colorectal cancer by annual colonoscopy or flexible sigmoidoscopy should be performed, beginning at puberty. In families with attenuated familial adenomatous polyposis (AFAP) or MAP, surveillance should be by colonoscopy (strong recommendation, moderate quality of evidence).
9. Absolute indications for immediate colectomy in FAP, AFAP, and MAP include: documented or suspected cancer or significant symptoms. Relative indications for surgery include the presence of multiple adenomas >6 mm, a significant increase in adenoma number, and inability to adequately survey the colon because of multiple diminutive polyps (strong recommendation, low quality of evidence).
10. Screening for gastric and proximal small bowel tumors should be done using upper endoscopy including duodenoscopy starting at age 25–30 years. Surveillance should be repeated every 0.5–4 years depending on Spigelman stage of duodenal polyposis: 0=4 years; I=2–3 years, II=1–3 years, III=6–12 months, and IV=surgical evaluation. Examination of the stomach should include random sampling of fundic gland polyps. Low-grade dysplasia is common in fundic gland polyps, and surgery should be reserved for high-grade dysplasia or cancer (strong recommendation, very low quality of evidence).
11. Annual thyroid screening by ultrasound should be recommended to individuals affected with FAP, MAP, and attenuated polyposis (conditional recommendation, low quality of evidence).
12. Biannual screening should be offered to affected infants until age 7 years with α -fetoprotein and ultrasounds (conditional recommendation, very low quality of evidence).
13. Postsurgical surveillance should include yearly endoscopy of rectum or ileal pouch, and examination of an ileostomy every 2 years (strong recommendation, low quality level of evidence).

*Hamartomatous polyposis syndromes**Peutz–Jeghers syndrome (PJS)*

14. Surveillance in affected or at-risk PJS patients should include monitoring for colon, stomach, small bowel, pancreas, breast, ovary, uterus, cervix, and testes cancers. Risk for lung cancer is increased, but no specific screening has been recommended. It would seem wise to consider annual chest radiograph or chest computed tomography (CT) in smokers (conditional recommendation, low quality of evidence).

Juvenile polyposis syndrome (JPS)

15. Surveillance of the gastrointestinal (GI) tract in affected or at-risk JPS patients should include screening for colon, stomach, and small bowel cancers (conditional recommendation, very low quality of evidence).
16. Colectomy and ileorectal anastomosis or proctocolectomy and ileal pouch-anal anastomosis is indicated for polyp-related symptoms, or when the polyps cannot be managed endoscopically (conditional recommendation, low quality of evidence).
17. Cardiovascular examination for and evaluation for hereditary hemorrhagic telangiectasia should be considered for *SMAD4* mutation carriers (conditional recommendation, very low quality of evidence).

Cowden syndrome (PTEN hamartoma tumor syndrome)

18. Surveillance in affected or at-risk Cowden syndrome patients should include screening for colon, stomach, small bowel, thyroid, breast, uterine, kidney, and skin (melanoma) cancers (conditional recommendation, low quality of evidence).

Serrated/hyperplastic polyposis syndrome

19. Patients with serrated polyposis should undergo colonoscopies every 1–3 years with attempted removal of all polyps >5 mm diameter (conditional recommendation, low quality of evidence).

20. Indications for surgery for serrated polyposis syndrome (SPS) include an inability to control the growth of serrated polyps, or the development of cancer. Colectomy and ileorectal anastomosis is a reasonable option given the risks of metachronous neoplasia (conditional recommendation, low quality of evidence).

21. There is no evidence to support extracolonic cancer surveillance for SPS at this time. Screening recommendations for family members are currently unclear pending further data and should be individualized based on results of baseline evaluations in family members (conditional recommendation, very low quality of evidence).

Hereditary pancreatic cancer

22. Surveillance of individuals with a genetic predisposition for pancreatic adenocarcinoma should ideally be performed in experienced centers utilizing a multidisciplinary approach and under research conditions. These individuals should be known mutation carriers from hereditary syndromes associated with increased risk of pancreatic cancer (Peutz–Jeghers, hereditary pancreatitis, familial atypical multiple melanoma and mole syndrome (FAMMM)) or members of familial pancreatic cancer kindreds with a pancreatic cancer affected first-degree relative. Because of a lower relative risk for pancreatic adenocarcinoma development in *BRCA1*, *BRCA2*, *PALB2*, *ATM*, and *LS* families, surveillance should be limited to mutation carriers with a first or second-degree relative affected with pancreatic cancer (conditional recommendation; very low quality of evidence).

23. Surveillance for pancreatic cancer should be with endoscopic ultrasound (EUS) and/or magnetic resonance imaging (MRI) of the pancreas annually starting at age 50 years, or 10 years younger than the earliest age of pancreatic cancer in the family. Patients with PJS should start surveillance at age 35 years (conditional recommendation, very low quality of evidence).

24. Because of the increased risk for pancreatic cancer development when compared with a pancreatic cyst in the sporadic setting, cystic lesion(s) of the pancreas detected during surveillance of a hereditary pancreatic cancer-prone family member requires evaluation by centers experienced in the care of these high-risk individuals. Determining when surgery is required for pancreatic lesions is difficult and is best individualized after multidisciplinary assessment (conditional recommendation, low quality of evidence).

Hereditary gastric cancer

Hereditary diffuse gastric cancer

25. Management for patients with hereditary diffuse gastric cancer should include: (i) prophylactic gastrectomy after age 20 years (>80% risk by age 80); (ii) breast cancer surveillance in women beginning at age 35 years with annual mammography and breast MRI and clinical breast examination every 6 months; and (iii) colonoscopy beginning at age 40 years for families that include colon cancer (conditional recommendation, low quality of evidence).

Table 3

GRADE (Grading of Recommendations Assessment, Development and Evaluation) system of evidence and strength of recommendation

High	Further research is very unlikely to change our confidence in the estimate of effect.
Moderate	Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.
Low	Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.
Very low	Any estimate of the effect is very uncertain.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 4**Amsterdam criteria, revised Bethesda guidelines, and colorectal cancer risk assessment tool***Amsterdam criteria I (24)*

At least three relatives with colorectal cancer (CRC); all of the following criteria should be present:

- One should be a first-degree relative of the other two;
- At least two successive generations must be affected;
- At least one of the relatives with CRC must have received the diagnosis before the age of 50 years;
- Familial adenomatous polyposis should be excluded;
- Tumors should be verified by pathologic examination.

Amsterdam criteria II (24)

At least three relatives must have a cancer associated with Lynch syndrome (colorectal, cancer of the endometrium, small bowel, ureter, or renal–pelvis); all of the following criteria should be present:

- One must be a first-degree relative of the other two;
- At least two successive generations must be affected;
- At least one relative with cancer associated with Lynch syndrome (LS) should be diagnosed before age 50;
- Familial adenomatous polyposis should be excluded in the CRC case(s) (if any);
- Tumors should be verified whenever possible.

Revised Bethesda guidelines (24)

Tumors from individuals should be tested for microsatellite instability (MSI) in the following situations:

- CRC diagnosed in a patient who is younger than 50 years of age
- Presence of synchronous, or metachronous, colorectal or other LS-related tumors^a, regardless of age
- CRC with MSI-high histology^b diagnosed in a patient who is younger than 60 years of age
- CRC diagnosed in a patient with one or more first-degree relatives with an LS-related cancer, with one of the cancers being diagnosed under age 50 years
- CRC diagnosed in a patient with two or more first- or second-degree relatives with LS-related cancer regardless of age

Colorectal cancer risk assessment tool^c

(Patient who answers yes to any question should have more comprehensive family history evaluation)

1. Do you have a first-degree relative (mother, father, brother, sister, or child) with any of the following conditions diagnosed before age 50?
 - Colon or rectal cancer
 - Cancer of the uterus, ovary, stomach, small intestine, urinary tract (kidney, ureter, bladder), bile ducts, pancreas, or brain
2. Have you had any of the following conditions diagnosed before age 50 years?
 - Colon or rectal cancer
 - Colon or rectal polyps
3. Do you have three or more relatives with a history of colon or rectal cancer?

(This includes parents, brothers, sisters, children, grandparents, aunts, uncles, and cousins)

^aLynch syndrome-related cancers include colorectal, endometrial, gastric, ovarian, pancreas, ureter and renal pelvis, biliary tract, brain (usually glioblastoma), small intestinal cancers, as well as sebaceous gland adenomas and keratoacanthomas.

^bPresence of tumor-infiltrating lymphocytes, Crohn's-like lymphocytic reaction, mucinous/signet-ring differentiation, or medullary growth pattern.

^cAdapted with permission from Kastrinos *et al.* (27).

Table 5

Cumulative risks of colorectal cancer in hereditary colorectal cancer syndromes

Syndrome	Gene	Risk	Average age of diagnosis (years)	References
Sporadic cancer		4.8%	69	SEER(303)
Lynch syndrome	<i>MLH1/MSH2</i>	M: 27–74% F: 22–61%	27–60	(30–35,38)
	<i>MSH6</i>	M: 22–69% F: 10–30% M/F: 12%	50–63	(31,36,49,64)
	<i>PMS2</i>	M: 20% F: 15%	47–66	(37)
Familial adenomatous polyposis (FAP)	<i>APC</i>	100%	38–41	(81,123,126,316)
Attenuated FAP	<i>APC</i>	69%	54–58	(88,90,126,317–319)
<i>MUTYH</i> -associated polyposis	<i>MUTYH</i>	43–100%	48–50	(109,126,134,135,319)
Juvenile polyposis	<i>SMAD4</i> <i>BMPRIA</i>	38–68%	34–44	(126,220)(320–323)
Peutz–Jeghers syndrome	<i>STK11</i>	39%	42–46	(126,196,197)
Cowden syndrome	<i>PTEN</i>	9–16%	44–48	(224,235,236,324)
Serrated polyposis syndrome	Not known	~>50%	48	(243,254)

F, female; M, male.

Table 6

Studies of colorectal screening in hereditary colorectal cancer (CRC) syndromes

Study (year)	Reference	Subjects	Design	Findings
<i>Lynch syndrome</i>				
Järvinen <i>et al.</i> (1995)	(42)	252 At-risk individuals from 20 of 22 families with MMR mutations.	Observational: all invited for colonoscopy screening; 133 had q 3-year colonoscopy, 118 declined.	62% Less CRC in screened ($P=0.03$). Tumor stage more favorable in screened. No deaths in screened vs. 5 deaths in nonscreened.
Järvinen <i>et al.</i> (2000)	(41)	252 At-risk individuals from 20 of 22 families with MMR mutations.	Observational: follow-up of Järvinen <i>et al.</i> (1995) study	62% reduction in CRC in screened ($P=0.02$). No deaths from CRC in screened vs. 9 deaths in nonscreened.
De Vos tot Nederveen Cappel <i>et al.</i> (2002)	(48)	857 Members of 114 HNPCC- or MMR-positive families.	Observational: tumor stage with more frequent (> 2 years) vs. less frequent colonoscopy; 10-year risk of CRC with partial vs. subtotal colectomy.	Earlier stage CRC with more frequent colonoscopy. 15.7% risk of CRC with partial vs. 3.4% with subtotal colectomy at 10 years.
Dove-Edwin <i>et al.</i> (2005)	(44)	554 At-risk members of 290 families with HNPCC or MMR mutations.	Prospective observational: evaluation of efficacy of colonoscopy.	Estimated 72% decrease in CRC death in screened individuals.
Jarvinen <i>et al.</i> (2009)	(43)	242 MMR mutation-positive and 367 mutation-negative subjects.	Observational: cancer incidence/survival of 11.5-year follow-up of colonoscopy surveillance	No increase in cancer mortality in mutation-positive vs. -negative individuals.
<i>Familial adenomatous polyposis (FAP)</i>				
Vasen <i>et al.</i> (1990)	(325)	230 Confirmed FAP cases	Observational: compared colorectal cancer rates in symptomatic cases vs. family members identified via screening	47% of the symptomatic cases had CRC at a mean age of 35 years compared with 4% at 24 years
Bjork <i>et al.</i> (2000)	(128)	195 Subjects with FAP	Observational: evaluation of rectal cancer morbidity and mortality after colectomy with ileorectal anastomosis (IRA)	The cumulative risk of rectal cancer 20 years after an IRA was 12.1%. There was a 7% cumulative risk of mortality.
Jarvinen (1992)	(82)	251 Affected subjects from 81 FAP families	Observational: evaluation of the effects of family screening on the occurrence of colorectal cancer by comparing cases diagnosed during family screening (call-up group) and symptomatic probands	65.5% Of probands had colorectal cancer compared with 6.6% of the call-up cases. The call-up group had a significantly increased lifetime cumulative survival from age 31 years.
<i>MUTYH-associated polyposis (MAP)</i>				
Nascimbeni <i>et al.</i> (2010)	(326)	14 <i>MUTYH</i> -positive subjects	Retrospective observational: to evaluate the risk of cancer or severe polyposis of the rectal stump after total colectomy	11 Cases underwent total colectomy with ileorectal anastomosis and yearly proctoscopic surveillance; no patient developed rectal cancer during surveillance (median duration: 5 years).

Study (year)	Reference	Subjects	Design	Findings
Nielsen <i>et al.</i> (2010)	(327)	147 Cases and 272 matched controls	Retrospective: compared survival between cases with MAP colorectal cancer and matched controls with colorectal cancer from the general population	Five-year survival was higher for the MAP colorectal subjects (78% vs. 63%). Survival remained better after adjusting for differences between the groups (hazard ration of 0.48).
<i>Peutz–Jeghers syndrome (PJS; see Table 8)</i>				
<i>Juvenile polyposis</i>				
Oncel <i>et al.</i> (2005)	(223)	13 Juvenile polyposis cases	Retrospective: to evaluate the long-term outcomes of the surgery in JPS cases who present with symptomatic colonic polyps	Eleven presented with rectal bleeding initially. Five of 10 patients who had initial rectum-preserving surgery required subsequent proctectomy; a total of 8 cases had their rectum removed during the study period. Following their ultimate operations, five patients required multiple polypectomies for recurrent pouch/rectal polyps.
<i>Cowden syndrome</i>				
Heald <i>et al.</i> (2010)	(224)	127 PTEN mutation carriers	Retrospective: to describe the gastrointestinal phenotype in a prospective series of PTEN mutation carriers.	Sixty-nine cases had 1 or more endoscopies and polyps were present in 64 (93%). Of the 64, half had hyperplastic polyps and 24 had both upper and lower gastrointestinal (GI) polyps; 9 (13%) cases had colorectal cancer at age <50 years.
<i>Sessile serrated polyposis syndrome</i>				
Boparai <i>et al.</i> (2010)	(259)	77 Hyperplastic polyposis syndrome cases	Retrospective: to describe the clinical and pathological features of a large hyperplastic polyposis syndrome (HPS) cohort	In a mean follow-up period of 5.6 years, 1,984 polyps were identified. Colorectal cancer was identified in 27 cases; 22 were diagnosed at initial colonoscopy. Four of the 5 interval CRCs were detected in diminutive serrated polyps (4–16 mm).

Table 7

Cumulative risks of extracolorectal cancer in hereditary colorectal cancer syndromes

Cancer site	General population risk ^a	Syndrome risk	Average age of diagnosis (years)	References
<i>Lynch syndrome</i>				
Endometrium				
General population	2.7%		65	SEER (328)
<i>MLH1/MSH2</i>		14–54%	48–54	(30,31,33,35,36,38)
<i>MSH6</i>		16–71%	53–54	(31,36,49)
<i>PMS2</i>		15%	49–50	(37)
Stomach	<1%	0.2–13%	49–55	(31,34,49,53–55,61,63,64)
Ovary	1.4%	3.4–22%	42–54	(31,36–38,49,53–55,63,64,67)
Hepatobiliary tract	<1%	0.02–4%	54–57 ^b	(31,55,63,67)
Urinary tract	<1%	0.2–25.5%	52–57	(31,37,49,53–55,63,64,67)
Small bowel	<1%	0.4–12%	46–51	(31,34,53–55,64)
Brain/central nervous system	<1%	1.2–3.7%	50–55	(53,55,63,64)
Sebaceous neoplasm	<1%	9% ^c	51–54	(329–331)
Pancreas	1.5%	0.4–3.7%	51.5–56.5 ^b	(33,53,271)
Prostate	15.3%	9–30%	59–60	(54,76,332)
Breast (female)	12.3%	1.5–18%	46–52	(53,54,75)
<i>Familial adenomatous polyposis(FAP)</i>				
Small bowel (duodenum/periampullary)	<1%	3–10%	44 50–52 ^b	(81,126,127,138,144,146,316,333)
Stomach	<1%	<1%	49	(126,138,316)
Pancreas	1.5%	1.7%	50 ^b	(126,138,334)
Thyroid	1.1%	2%	25–33	(126,138,171,172,316,334,335)
Liver (hepatoblastoma)	<1%	1–2%	Most often occurs in the first 5 years of life	(126,138,173)
Brain/central nervous system	<1%	1–2%	15–21	(126,138,336–338)
<i>Attenuated FAP</i>				
Small bowel (duodenum/periampullary)	<1%	4–12%	60	(90,126)
Thyroid	1.1%	1–2%	26	(90,126)
<i>MUTYH-associated polyposis</i>				
Small bowel (duodenum)	<1%	4%	61 ^b	(126,174)
Stomach	<1%	1%	38 ^b	(126,174)
<i>Juvenile polyposis</i>				
Upper gastrointestinal (GI) cancer (stomach, pancreas, and small bowel)	—	21% ^c	54	(126,183,321)
<i>Peutz–Jeghers syndrome</i>				

Cancer site	General population risk ^a	Syndrome risk	Average age of diagnosis (years)	References
Stomach	<1%	29%	30–40	(196,197)
Small bowel	<1%	13%	37–42	(196,197)
Pancreas	1.5%	11–36%	41–52	(194,196,197,339)
Breast	12.4%	32–54%	37–59	(194,196,197)
Ovarian (mostly SCTAT (sex cord tumor with annular tubules))	1.6%	21%	28	(197)
Uterus	2.7%	9%	43	(196,197)
Cervix (adenoma malignum)	<1%	10%	34–40	(197,340)
Testicular (Sertoli cell tumor)	<1%	9%	6–9	(196,197)
Lung	6.9%	7–17%	47	(194,196,197)
<i>Cowden syndrome</i>				
Breast	12.4%	25–85%	38–46	(184,235,236,341–343)
Thyroid	1.1%	3–38%	31–38 ^b	(184,235,236,342–345)
Endometrium	2.7%	5–28%	25 ^d	(184,228,235,236)
Kidney (renal cell)	1.6%	15–34%	40 ^d	(183,224,225,235,346,347)
Melanoma	2	6%	3 ^e	(183,224,225,235,346,347)

^a SEER, 2013 (328).

^b Median age.

^c Incidence.

^d Onset of risk.

^e Youngest age of onset.

Table 8

Studies of extracolonic cancer screening in hereditary colorectal cancer syndromes

Organ	Study (years)	Reference	Subjects	Design	Findings
<i>Lynch syndrome (LS)</i>					
Endometrial and ovarian	Dove-Edwin <i>et al.</i> (2002)	(56)	292 Women from hereditary nonpolyposis colorectal cancer (HNPCC) or HNPCC-like families.	Observational: all offered transvaginal ultrasounds.	Two cases of endometrial cancer (EC) presented with symptoms, neither detected by ultrasound.
	Rijcken <i>et al.</i> (2003)	(348)	41 Women with MMR mutations or fulfilled Amsterdam I criteria followed for median of 5 years.	Observational: all offered annual pelvic exam, transvaginal ultrasound, CA-125.	17 Of 179 ultrasounds gave reason for endometrial sampling with 3 premalignant lesions noted; one interval endometrial cancer presented symptomatically.
	Renkonen-Sinisalo <i>et al.</i> (2007)	(57)	175 Women with MMR mutations.	Observational: all offered transvaginal ultrasound and endometrial biopsy.	14 cases of EC; 11 diagnosed by surveillance. Biopsy diagnosed 8 of 11 EC and 14 cases of premalignant hyperplasia. Ultrasound indicated 4 EC cases but missed 6 others. 4 case of ovarian cancer, none found by ultrasound.
	Lecuru <i>et al.</i> (2008)	(349)	62 Women (13 with MMR mutation, 49 met Amsterdam II criteria).	Observational: annual hysteroscopy and endometrial biopsy.	3 Malignancies in 3 patients with abnormal bleeding; 3 cases of hyperplasia in asymptomatic patients; hysteroscopy 100% sensitive for cancer or hyperplasia.
	Gerritzen <i>et al.</i> (2009)	(350)	100 Women from families with MMR mutation.	Observational: annual transvaginal ultrasound, CA-125, endometrial sampling.	3 Atypical hyperplasias and 1 endometrial cancer diagnosed. One stage III ovarian cancer developed despite ultrasound.
	Stuckless <i>et al.</i> (2013)	(351)	174 Women with MSH2 gene mutation.	Case-control: Cases: 54 patient with at least one screening exam (transvaginal, endometrial biopsy, or CA-125 test); Controls: matched women without screening.	Stage I/II cancer diagnosed in 92% of screened patients compared with 71% in control group ($P=0.17$). Two of three deaths in the screened group from ovarian cancer.
	Stuckless <i>et al.</i> (2012)	(46)	322 MSH2 mutation carriers.	Observational: cancer incidence and survival in 152 screened vs. 170 not screened by colonoscopy.	Median age to CRC later in screened vs. nonscreened. Survival statistically improved in screened vs. nonscreened.
Gastric	Schmeler <i>et al.</i> (2006)	(58)	315 Women with MMR mutation with and without gynecological surgery.	Retrospective: risk of uterine and ovarian cancer in patients with and without prophylactic/clinically indicated gynecological surgery.	No uterine or ovarian cancer in surgery group vs. 33 and 5% cancer respectively in nonsurgery group.
	Renkonen-Sinisalo <i>et al.</i> (2002)	(352)	73 Patients with MMR mutation; 32 MMR	Observational: Upper endoscopy with gastric biopsies.	In MMR gene-positive patients, <i>H. pylori</i> in 26%, atrophy 14%, intestinal metaplasia 14%. No statistical

Organ	Study (years)	Reference	Subjects	Design	Findings
			mutation-negative family members.		difference between gene-positive and -negative groups.
Small bowel	Saurin <i>et al.</i> (2010)	(66)	35 Patients with MMR mutations.	Observational: Capsule endoscopy and computed tomographic (CT) enteroclysis screening of small bowel.	Small bowel neoplasms were found in 8.6% of cases (1 patient with jejunal carcinoma and 2 with jejunal adenoma). Capsule endoscopy found all lesions; CT enteroscopy found cancer but missed adenomas.
Urinary tract	Myrhoj <i>et al.</i> (2008)	(68)	977 At-risk individuals in families suspected to have HNPCC/LS.	Observational: Retrospective review of screening urine cytology (UC) and diagnosis of urinary cancer.	0.1% Of UC exams lead to diagnosis of urothelial tumor. 10× more UC exams lead to false-positive diagnosis. Sensitivity of UC was 29%.
<i>Familial adenomatous polyposis (FAP)</i>					
Duodenum	Bulow <i>et al.</i> (2004)	(144)	367 FAP cases	Prospective: to describe the long-term natural history of duodenal adenomatosis in FAP and evaluate the indications for prophylactic duodenal surveillance	65% Of cases had duodenal adenomas on their first endoscopy. The cumulative incidence of duodenal carcinoma was 4.5% at 57 years; cases with Spigelman stage IV adenomatosis on initial endoscopy were at higher risk compared with those with stages 0–III
	Biasco <i>et al.</i> (2006)	(353)	50 FAP cases	Prospective: to evaluate the presence and severity of precancerous duodenal mucosal lesions	86% Of cases had duodenal lesions at the end of the study compared with 38% at first endoscopy. Eleven subjects developed stage IV disease and had endoscopic or surgical resection.
Thyroid	Jarrar <i>et al.</i> (2011)	(354)	192 FAP cases	Prospective: to clarify the incidence of thyroid cancer in patients with FAP	38% Of cases had thyroid nodules and 2.6% had thyroid cancer. Clinical history and neck examination did not detect any of the cancers
<i>Peutz–Jeghers syndrome (PJS)</i>					
Gastrointestinal (GI) tract	Latchford <i>et al.</i> (2011)	(355)	63 PJS cases from 48 pedigrees	Retrospective review: to assess outcomes from GI surveillance in patients with PJS	Baseline investigations were done in 12 subjects. The rest of the patients were followed for a median of 10 years and 776 procedures were performed. Of the 2,461 polypectomies done, 6 polyps contained atypia or dysplasia; there were two cases of perforation following resection of polyps >2 cm. No luminal GI cancers were diagnosed.
Small bowel	Brown <i>et al.</i> (2006)	(356)	19 Adult PJS cases	Prospective: to evaluate the performance of capsule endoscopy in small bowel surveillance of adults with PJS vs. barium follow-through	Capsule endoscopy detected more significant polyps than barium follow-through but seemed less reliable for accurately sizing 1–2 cm polyps.
	Gupta <i>et al.</i> (2010)	(357)	19 Adult PJS cases	Prospective: to assess the utility of magnetic resonance (MR) enterography compared with capsule endoscopy for small bowel	All cases underwent both procedures. MR enterography detected large polyps (>15 mm) missed in three patients by capsule endoscopy. Size assessments of large polyps

Organ	Study (years)	Reference	Subjects	Design	Findings
Pancreas	Poley <i>et al.</i> (2009)	(358)	44 Individuals at high risk of developing pancreatic cancer (2 PJS cases)	polyp detection in PJS Prospective: to investigate the use of endoscopic ultrasound for screening individuals at high risk of developing pancreatic cancer	appeared more reproducible with MR enterography. Initial screening detected an asymptomatic mass lesion in 3 cases (6.8%) and premalignant intraductal papillary mucinous neoplasm (IPMN)-like lesions in 16%.
<i>Cowden syndrome</i> Thyroid	Milas <i>et al.</i> (2012)	(345)	225 PTEN mutative-positive cases	Retrospective: to characterize Cowden syndrome-associated malignant and benign thyroid disease	32 Cases (14%) had thyroid cancer (mostly papillary type) at a median age of 35 years. Initial thyroid ultrasound in 16 of 25 subjects revealed thyroiditis/goiters in all >13 years of age.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 9Duodenal adenomatosis staging system^a

Polyps	1 Point	2 Points	3 Points
Number	<4	5–20	>20
Size	0–4 mm	5–10 mm	>10
Histology	Tubular	Tubulovillous	Villous
Dysplasia	Mild	Moderate	Severe

Spigelman stage	Total points	Frequency of surveillance
		Recommended duodenal surveillance frequency^b
0	0	Every 4 years
I	4	Every 2–3 years
II	5–6	Every 1–3 years
III	7–8	Every 6–12 months
IV	9–12	Expert surveillance every 3–6 months Surgical evaluation
Complete mucosectomy or duodenectomy or Whipple procedure if duodenal papilla is involved		

^a Adapted from ref. (154).

^b Adapted with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines) for Genetic/Familial High-Risk Assessment: Colorectal V.1.2014. 2014 National Comprehensive Cancer Network (24) The NCCN Guidelines and illustrations herein may not be reproduced in any form for any purpose without the express written permission of the NCCN. To view the most recent and complete version of the NCCN Guidelines, go online to NCCN.org. NATIONAL COMPREHENSIVE CANCER NETWORK, NCCN, NCCN GUIDELINES, and all other NCCN content are trademarks owned by the National Comprehensive Cancer Network.

Table 10

Surveillance recommendations for hereditary gastrointestinal (GI) cancer syndromes

Site	Age to begin surveillance (years)	Surveillance interval (years)	Surveillance procedures and comments	References
<i>Lynch syndrome</i>				
Colon	20–25 (<i>MLH1/MSH2</i> mutation) 25–30 (<i>MSH6/PMS2</i> mutation)	1–2 ^a	Colonoscopy ^a	(24,29,47)
Endometrial and ovarian	30–35	1	No evidence of survival benefit of surveillance proven (see Table 8). Annual pelvic exam; offer annual endometrial biopsy and transvaginal ultrasound to at-risk women. Consider total abdominal hysterectomy/bilateral salpingo-oophorectomy in women who have completed childbearing.	(24,58,60,359)
Pancreas		1–2	Consider magnetic resonance imaging (MRI) and/or endoscopic ultrasound in mismatch repair gene mutation carriers with pancreas cancer in a first-degree relative	(73)
Urinary tract	25–30	1	Limited data exist to advocate urinary screening (NCCN recommends considering an annual urinalysis).	(24)
Small bowel and gastric	30–35	3–5	Consider esophagogastroduodenoscopy with extended duodenoscopy in select individuals	(24)
<i>Familial adenomatous polyposis (FAP)</i>				
Colon	10–15	1–2 ^b	Flexible sigmoidoscopy or colonoscopy ^b	(24,127)
Upper gastrointestinal	25–30	1–5 ^c	Esophagogastroduodenoscopy with a side-viewing instrument	(127)
Thyroid	Late teenage years	1	Annual thyroid examination; annual thyroid ultrasound	(24,171)
Intraabdominal desmoids		1	Annual abdominal palpation. (NCCN Guidelines: Consider abdominal MRI or computed tomography (CT) 1–3 years after colectomy then at 5–10-year intervals with family history of symptomatic desmoids or if suggestive abdominal symptoms occur).	(24)
<i>Attenuated FAP</i>				
Colon	18–20	1–2 ^b	Colonoscopy	(24,127)
Upper gastrointestinal	25–30	1–5 ^c	Esophagogastroduodenoscopy with a side-viewing instrument	(127)
Thyroid		1	Annual thyroid examination	(24)
<i>MUTYH-associated polyposis</i>				
Colon	25–30	1–2 ^b	Colonoscopy	(127)

Site	Age to begin surveillance (years)	Surveillance interval (years)	Surveillance procedures and comments	References
Upper gastrointestinal	30–35	Baseline ^c	Esophagogastroduodenoscopy with a side-viewing instrument	(24)
<i>Peutz–Jeghers syndrome</i>				
Colon	8, 18 ^d	3	Colonoscopy ^d	(182)
Stomach	8, 18 ^d	3	Esophagogastroduodenoscopy ^d	(182)
Small bowel	8, 18 ^d	3	Video capsule endoscopy ^d	(182)
Pancreas	30	1–2	Magnetic resonance cholangiopancreatography or endoscopic ultrasound	(24,182,196)
Breast	25	1	Annual self-exam starting age 18, annual breast MRI, and/or mammogram starting at age 25	(182,196,360)
Ovarian	25	1	Pelvic exam and pelvic or transvaginal ultrasound, CA-125 probably not helpful	(182)
Endometrial	25	1	Pelvic exam and pelvic or transvaginal ultrasound	(182)
Cervix (adenoma malignum)	25	1	Pap smear	(182,196)
SCTAT (sex cord tumor with annular tubules)	25	1	Same as uterine and ovarian; almost all women develop SCTAT, but 20% become malignant	(182)
Testicular (Sertoli cell tumor)	Birth to teenage years	1	Testicular exam, ultrasound if abnormalities palpated or if feminization occurs; 10 to 20% of benign Sertoli cell tumors become malignant	(182)
Lung	—	—	Provide education about symptoms and smoking cessation	(24)
<i>Juvenile polyposis syndrome</i>				
Colon	12–15	1–3	Colonoscopy ^e	(24,210)
Stomach	12–15	1–3	Esophagogastroduodenoscopy ^e	(24,208)
Small Intestine	—	—	Rare, undefined lifetime risk. Periodic enteroscopy, capsule endoscopy, and/or CT enterography	(24,208)
Pancreas	—	—	Rare, undefined lifetime risk. No screening recommendations given	(24)
HHT (hereditary hemorrhagic telangiectasia)	Within first 6 months of life	—	Undefined lifetime risk. In individuals with SMAD4 mutations, screen for vascular lesions associated with HHT	(24,361)
<i>Cowden syndrome^f</i>				
Colon	15	2	Colonoscopy, intervals may increase or decrease, depending on findings	(183)
Upper GI tract and small bowel	15	2–3	Esophagogastroduodenoscopy. If duodenal polyposis is present, repeat depending on number of polyps	(183)

Site	Age to begin surveillance (years)	Surveillance interval (years)	Surveillance procedures and comments	References
Thyroid	Adolescence	1	Thyroid exam and baseline ultrasound	(183,347)
Breast	25 30–35	Monthly 1	Self-breast exam Mammography and breast magnetic resonance imaging	(24)
Uterine	30–35	1	Annual endometrial sampling or vaginal ultrasound	(24)
Renal cell	18	1	Urine analysis with cytology and possibly renal ultrasound	(235)
Melanoma	By 18	1	Physical cutaneous examination	(235)
<i>Serrated polyposis syndrome</i>				
Colon	?	1–3	Colonoscopy ^g	(24)
<i>Hereditary pancreatic cancer</i>				
Pancreas	50 ^h	1 ⁱ	Endoscopic ultrasound and/or MRI/magnetic resonance cholangiopancreatography (MRCP) ⁱ Provide education about smoking cessation	(295,302)
<i>Hereditary gastric cancer</i>				
Stomach	5–10 Years before earliest cancer in family	0.5–1	Esophagogastroduodenoscopy; detailed 30-min exam with multiple random biopsies	(313,314)
Breast	35	1	Breast exam, annual mammography Annual breast magnetic resonance imaging	
Colon	40	?	Colonoscopy ^j	

^a Consider annual colonoscopy in confirmed mutation carriers.

^b Annual colonoscopy if surgery is delayed for >1 year after onset of colon polyps.

^c Frequency depends on severity of duodenal polyposis.

^d Start at age 8 years; if polyps present, repeat every 3 years; if no polyps, repeat at age 18, then every 3 years, or earlier if symptoms occur.

^e Start at age 12 years; if polyps present, repeat annually; if no polyps, repeat every 2–3 years.

^f Baseline physical examination at diagnosis and annual targeted history and physical examination thereafter. (235)

^g Colonoscopy with polypectomy until all polyps > 5 mm are removed, then colonoscopy every 1–3 years.

^h Or 10 years younger than the earliest age of pancreatic cancer in the family.

ⁱ MRI/MRCP. Frequency depends on severity of observed lesions and the need for pathologic examination.

^j Consider in hereditary gastric cancer families with individuals affected by colon cancer.

Table 11Indications for genetic evaluation for Cowden syndrome (PTEN hamartoma tumor syndrome)^a

Individual from a family with a known *PTEN* mutation

Individual meeting clinical diagnostic criteria for Cowden syndrome (CS) as outlined below

Individual with a personal history of any of the following:

- Bannayan–Riley–Ruvalcaba syndrome (BRRS)
- Adult Lhermitte–Duclos disease
- Autism spectrum disorder and macrocephaly
- Two or more biopsy-proven trichilemmomas
- Two or more major criteria (one must be macrocephaly)
- Three major criteria, without macrocephaly
- One major and three minor criteria^b
- Four minor criteria

At-risk individual with one major or two minor criteria and a relative with a clinical diagnosis of CS or BRRS for whom testing has not been performed

Major criteria

- Breast cancer
- Endometrial cancer
- Follicular thyroid cancer
- Multiple gastrointestinal hamartomas or ganglioneuromas
- Macrocephaly (megaloccephaly, 97th percentile)
- Macular pigmentation of glans penis
- Mucocutaneous lesions alone if:
 - One biopsy proven trichilemmoma, or
 - Multiple palmoplantar keratoses, or
 - Multifocal or extensive oral mucosal papillomatosis, or
 - Multiple cutaneous facial papules (often verrucous)

Minor criteria

- Autism spectrum disorder
- Colon cancer
- Esophageal glycogenic acanthosis (3)
- Lipomas
- Mental retardation (i.e., IQ < 75)
- Papillary or follicular variant of papillary thyroid cancer
- Thyroid structural lesions (e.g., adenoma, nodule(s), goiter)
- Renal cell carcinoma
- Single gastrointestinal hamartoma or ganglioneuroma
- Testicular lipomatosis
- Vascular anomalies (including multiple intracranial developmental venous anomalies)

^a Adapted with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines) for Genetic/Familial High-Risk Assessment: Breast and Ovarian V.4.2013.(3) 2014 National Comprehensive Cancer Network. All rights reserved. The NCCN Guidelines and illustrations herein may not be reproduced in any form for any purpose without the express written permission of the NCCN. To view the most

recent and complete version of the NCCN Guidelines, go online to NCCN.org. NATIONAL COMPREHENSIVE CANCER NETWORK, NCCN, NCCN GUIDELINES, and all other NCCN content are trademarks owned by the National Comprehensive Cancer Network.

^b If an individual has two or more major criteria, such as breast cancer and nonmedullary thyroid cancer, but does not have macrocephaly, one of the major criteria may be included as one of the three minor criteria to meet testing criteria.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 12

Syndromes associated with pancreatic adenocarcinoma and risk of developing pancreatic cancer

Syndrome	Relative risk of pancreatic cancer	Gene	References
Familial atypical multiple melanoma and mole (FAMMM)	13- to 39-fold	<i>CDKN2A</i>	(265–267)
Familial breast and ovarian	2-fold 3- to 9-fold	<i>BRCA1 BRCA2</i>	(268)(269,270)
Fanconi anemia, breast CA	Unknown	<i>PALB2</i>	
Familial adenomatous polyposis (FAP)	5-fold	<i>APC</i>	(334)
Lynch	9- to 11-fold	<i>MLH1, MSH2MSH6, PMS2</i>	(71,362)
Peutz–Jeghers syndrome	Up to 132-fold	<i>STK11/LKB1</i>	(197)
Li–Fraumeni	Unknown	<i>p53</i>	
Hereditary pancreatitis	53-fold	<i>PRSS1</i>	(264)
Ataxia–telangiectasia	3-fold	<i>ATM</i>	(272)
Familial pancreatic cancer:			
1 Or 2 first-degree relatives	4- to 7-fold	See Table 13	(276,277)
3 First-degree relatives	17- to 32-fold	Majority unknown	

Table 13

Yield for germline mutations In familial pancreatic cancer (FPC) kindreds without criteria for known genetic syndromes^a

Mutation	Yield of testing	References
<i>BRCA1</i>	0 to 6%	(281,286)
<i>BRCA2</i>	0 to 6%	(281,287,288)
<i>CDKN2A</i>	0 to 20%	(280,281,288)
<i>PALB2</i>	0 to 5%	(281,285,288)
<i>ATM</i>	2.4%	(283)
<i>STK11</i>	0%	(288)
<i>PALLD</i>	0%	(289,363)

^aUnknown for Lynch syndrome, familial adenomatous polyposis (FAP), and Li-Fraumeni syndrome.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 14**Standards for informed consent for genetic testing in gastrointestinal (GI) practice (364)**

Components of a proper informed consent for cancer genetic testing should include:

1. Information on the specific genetic mutation(s) or genomic variant(s) being tested, including whether or not the range of risk associated with the variant will affect medical care.
 2. Implications of positive and negative results.
 3. Possibility that the test will not be informative.
 4. Options for risk estimation without genetic or genomic testing.
 5. Risk of passing a genetic variant to children.
 6. Technical accuracy of the test, including, where required by law, licensure of the testing laboratory
 7. Fees involved in testing and counseling and, for direct to consumer testing, whether the counselor is employed by the testing company.
 8. Psychological implications of test results (benefits and risks).
 9. Risks and protections against genetic discrimination by employers or insurers.
 10. Confidentiality issues, including, for direct-to-consumer testing companies, policies related to privacy and data security.
 11. Possible use of DNA testing samples in future research.
 12. Options and limitations of medical surveillance and strategies for prevention after genetic or genomic testing.
 13. Importance of sharing genetic and genomic test results with at-risk relatives so that they may benefit from this information.
 14. Plans for follow-up after testing.
-

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript